

Adapting Stable Matchings to Evolving Preferences*

Robert Bredereck¹, Jiehua Chen^{†2}, Dušan Knop^{‡3}, Junjie Luo^{§1}, and Rolf Niedermeier¹

¹Technische Universität Berlin, Chair of Algorithmics and Computational Complexity
`{robert.bredereck, junjie.luo, rolf.niedermeier}@tu-berlin.de`

²Algorithms and Complexity Group, TU Wien, Vienna, Austria `jiehua.chen2@gmail.com`

³Department of Theoretical Computer Science, Faculty of Information Technology, Czech Technical University in Prague, Prague, Czech Republic `dušan.knop@fit.cvut.cz`

November 25, 2019

Abstract

Adaptivity to changing environments and constraints is key to success in modern society. We address this by proposing “incrementalized versions” of STABLE MARRIAGE and STABLE ROOMMATES. That is, we try to answer the following question: for both problems, what is the computational cost of adapting an existing stable matching after some of the preferences of the agents have changed. While doing so, we also model the constraint that the new stable matching shall be not too different from the old one. After formalizing these incremental versions, we provide a fairly comprehensive picture of the computational complexity landscape of INCREMENTAL STABLE MARRIAGE and INCREMENTAL STABLE ROOMMATES. To this end, we exploit the parameters “degree of change” both in the input (difference between old and new preference profile) and in the output (difference between old and new stable matching). We obtain both hardness and tractability results, in particular showing a fixed-parameter tractability result with respect to the parameter “distance between old and new stable matching”.

1 Introduction

Imagine the following scenario. A manager responsible for a group of $2n$ workers has to form n two-worker teams based on the preferences over potential work partners of each worker. The manager, being interested in a robust solution, computes a stable matching (indeed, this refers to the STABLE ROOMMATES problem). However, say every month the workers may update their preferences about wanted work partners (the updates may be based on gained experiences, new information, newly developed personal skills etc.). The team manager then has to find a new stable matching respecting the individually evolved preferences. To this end, however, the team manager may not want to allow too radical changes in the composition of the two-worker teams because this might e.g. overburden administration. Thus, a moderate change is acceptable, but too radical changes in the team compositions should be avoided whenever possible. We address this scenario by introducing and studying “incremental versions” of the two most prominent stable matching problems, namely STABLE MARRIAGE and STABLE ROOMMATES.¹

*We thank anonymous reviewers of AAAI for their insightful comments.

[†]Main work done while JC was with University of Warsaw, where she was supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme under grant agreement numbers 677651. JC was also supported by the WWTF research project (VRG18-012).

[‡]Main work done while DK was with TU Berlin, supported by the DFG project MaMu (NI 369/19).

[§]Supported by CAS-DAAD Joint Fellowship Program for Doctoral Students of UCAS and by the DFG project AFFA (BR 5207/1 and NI 369/15)

¹The naming “Incremental” is inspired by work in the context of clustering and information retrieval [6].

In stable matching scenarios or, in other words, in matching under preferences [28], one is given a set of agents, each of them having preferences over (some of) the other agents, and the goal is to match pairs of agents such that the outcome is stable. Informally speaking, stability means that there are no two agents that would both prefer to be matched with each other instead to their current partners in the matching (or being unmatched). Two classic problems here are the bipartite case with two equal-sized sets of agents (referred to as STABLE MARRIAGE) and the general case of an even number of agents (referred to as STABLE ROOMMATES). Motivated by the introductory considerations, we next define “incremental versions” of both problems; further formal definitions are presented later.

INCREMENTAL STABLE MARRIAGE

Input: Two disjoint sets U and W of n agents each, two preference profiles \mathcal{P}_1 and \mathcal{P}_2 for $U \uplus W$, a stable matching M_1 for profile \mathcal{P}_1 , and a non-negative integer k .

Question: Does $U \uplus W$ admit a stable matching M_2 for profile \mathcal{P}_2 such that $\text{dist}(M_1, M_2) = |M_1 \Delta M_2| \leq k$?

Herein, $M_1 \Delta M_2$ denotes the symmetric difference between sets M_1 and M_2 . The incremental setting for STABLE ROOMMATES is defined analogously.

INCREMENTAL STABLE ROOMMATES

Input: A set V of $2n$ agents, two preference profiles \mathcal{P}_1 and \mathcal{P}_2 for V , a stable matching M_1 for profile \mathcal{P}_1 , and a non-negative integer k .

Question: Does V admit a stable matching M_2 for profile \mathcal{P}_2 such that $\text{dist}(M_1, M_2) = |M_1 \Delta M_2| \leq k$?

In both definitions there are two main regulating screws. First, we are given *two* preference profiles, the old \mathcal{P}_1 and the new \mathcal{P}_2 , thereby reflecting the change of preferences. Indeed, to reflect only moderate changes (“evolution”), we will subsequently measure the difference between the two profiles (later referred to as swap distance), yielding a natural problem-specific parameter (the smaller it is, the less revolutionary the changes are). Second, the number k can be interpreted as a locality parameter—it exposes how close the new matching has to be to the old one. The smaller we choose k , the more conservative we are with respect to change in the outcome (namely the difference between old and new stable matching). Together, we thus have one parameter to regulate the degree of change measured in the *input* preferences and one parameter to regulate the degree of change measured in the *output* stable matching solution. Taking up these two parameters, we provide a thorough parameterized complexity analysis of these kinds of stable matching problems with evolving preferences.² To this end, we also distinguish between preferences with and without ties. Roughly speaking, we provide positive (fixed-parameter) tractability results in the case without ties and several (parameterized) intractability results for the case with ties. Before describing our results in more detail, however, we discuss related work.

1.1 Related work

There is previous work on matching-related problems in the context of dynamic graph algorithms where vertices and/or edges arrive or depart iteratively over time. The goal then is to maintain a solution of sufficient quality by performing necessary updates after every single change. The main difference to our work is that we study changes between two preference profiles (not so

²In the large field of dynamic graph algorithms, which significantly differs from our setting, the term “dynamic” refers to more fine-grained scenarios where typically edges and/or vertices may be added or deleted in a stepwise fashion, and one wants to efficiently update a solution after every such single change. In particular, this has also been studied in the (popular and stable) matching context [3, 17, 32]. The main difference to our work is that we study changes between two preference profiles (not so much in the graph structure), and the changes can be at a larger scale. Moreover, we perform parameterized complexity studies which do not play a role in this previous work.

much in the graph structure), and the changes can be at a larger scale. For instance, Bhattacharya et al. [3] studied the maintenance of near-popular matchings (a scenario related to stable matchings) based on a greedy improvement strategy, heavily employing maximum vertex degree in their analysis. Ghosal et al. [17] and Nimbhorkar and Rameshwar [32] investigated dynamic rank-maximal and popular matchings, again within the setting of dynamic graph algorithms and a focus on update times after each single change in the graph. Kanade et al. [23] studied STABLE MARRIAGE where at each time-step, two random adjacent agents in some preference list are swapped, and designed approximation algorithms to maintain a matching with logarithmic number of blocking pairs.

Genc et al. [13, 14, 15, 16] studied *robustness* of a matching in STABLE MARRIAGE. Herein, the robustness of a given stable matching is measured by the number of modifications needed to find an alternative stable matching if some currently matched agent pairs break up. They define an (x, y) -supermatch as a stable matching that satisfies the following: If any x agents break up, then it is possible to rematch these x agents so that the new matching is again stable; further, this re-matching *must not* break up more than y other pairs. Genc et al. [16] showed that deciding whether a $(1, 1)$ -supermatch exists is NP-complete. The main distinguishing features compared to our model are that, using our two regulating screws mentioned above, we can model both moderate changes in the preferences (that is, the input) and moderate differences between the old and the new stable matching (that is, the output). Indeed, we perform a parameterized complexity analysis exploiting these parameters while Genc et al. [13, 14, 15, 16] focused on classic complexity results and used heuristics in experimental work. Moreover, we model changing preferences while they addressed breaking up matched pairs. Finally, our main contributions are in the STABLE ROOMMATES case while they exclusively focused on STABLE MARRIAGE without ties.

Our model of measuring the distance between the input and sought matching is related to the stable matching problem with *forbidden* and *forced* edges studied by Cseh and Manlove [10]. The goal is to find a stable matching which minimizes the number of forbidden edges plus the number of non-forced edges. While we do not model forbidden edges, Cseh and Manlove [10] do not assume changes in agents' preferences. This makes a difference in terms of parameterized complexity.

Marx and Schlotter [29, 30] studied local search aspects for the NP-hard STABLE MARRIAGE with ties. More specifically, they investigated the parameterized complexity of a local search variant of STABLE MARRIAGE using parameters such as the number of ties. Thus, the main overlap with our work is in terms of searching for local improvements and employing parameterized complexity analysis—the studied computational problems are different from ours as they do not model changes in the input preferences.

There have been numerous other models and investigations to enrich the basic stable matching model, including the use of only partially ordered preferences [11], “multilayer” stable matchings with several preference profiles to be obeyed ‘in parallel’ [2, 8, 31], or studying robust stability in a probabilistic model [26, 27] or from a quantitative angle [9].

Finally, let us briefly mention that the motivation for our incremental scenario for stable matching is related to similar scenarios in the context of clustering [6, 25], coloring [20], other dynamic versions of parameterized problems [1, 24], and reoptimization [4, 33].

1.2 Our contributions

Besides introducing a fresh model of stable matching computations, we provide results mostly in terms of parameterized complexity analysis for both INCREMENTAL STABLE MARRIAGE and INCREMENTAL STABLE ROOMMATES, where we see the main technical contributions mostly for the latter. In particular, our main algorithmic result is that INCREMENTAL STABLE ROOMMATES for input instances without ties is fixed-parameter tractable with respect to the parameter k (distance between the old and the new matchings). To show this, we heavily use structural

Ties	$ \mathcal{P}_1 \oplus \mathcal{P}_2 $	INCR. STABLE MARRIAGE		INCR. STABLE ROOMMATES	
no	any	P	(Prop. 2)	FPT for $k := \text{dist}(M_1, M_2)$ W[1]-hard for $k' := M_1 \cap M_2 $	(Thm. 1) (Prop. 6)
yes	1	W[1]-hard for k (Thm. 2)		W[1]-hard for k , even for compl. pref. NP-hard even if $k' \geq 0$	(Thm. 4) (Thm. 4)
yes	2	W[1]-hard for k' (Thm. 3)		W[1]-hard for k'	(Thm. 3)
yes	any	XP for k (Prop. 3)		XP for k	(Prop. 3)

Table 1: Overview of our results. Unless otherwise stated, results are for the general case where preferences could be incomplete.

results due to Irving [22] and Gusfield [18] and show how to exploit them for designing a fixed-parameter algorithm.

Most of our results are surveyed in Table 1. Herein, $\mathcal{P}_1 \oplus \mathcal{P}_2$ denotes the swap distance between two preference profiles (see Section 2 for formal definitions). The table indicates that we obtained a fairly complete picture of the computational complexity landscape, e.g., also complementing W[1]-hardness results with corresponding XP-algorithms.

Finally, we mention in passing that to achieve our results, throughout the work we also introduce and study some intermediate problems (for instance, EDGE-INCREMENTAL INDEPENDENT SET) which may be of independent interest and prove useful in other settings.

Organization of the Paper. The rest of the paper is organized as follows. We start with formal definitions and notations in Section 2. Our main algorithmic result Theorem 1 and other algorithms are presented in Section 3. Our hardness results are given in Section 4. We conclude in Section 5 with some open problems.

2 Definitions and notations

In this section, we review fundamental concepts used in matchings under preferences.

Preference lists and profiles. Let $V = \{1, 2, \dots, 2n\}$ be a set of $2n$ agents. Each agent $i \in V$ has a subset $V_i \subseteq V \setminus \{i\}$ of agents which they find *acceptable* as partners and has a *preference list* \succeq_i on V_i (i.e., a transitive and complete binary relation on V_i). Here, $x \succeq_i y$ means that i weakly prefers x over y (i.e. x is at least as good as y). We use \succ_i to denote the asymmetric part (i.e., $x \succeq_i y$ and $\neg(y \succeq_i x)$) and \sim_i to denote the symmetric part of \succeq_i (i.e., $x \succeq_i y$ and $y \succeq_i x$). If $x \sim_i y$, then we also say that x and y are *tied* in i 's preference list and that agent i has *ties* in their preference list.

For two agents x and y , we call x *most acceptable* to y if x is a maximal element in the preference list of y . Note that an agent can have more than one most acceptable agent. For two disjoint subsets of agents $X \subseteq V$ and $Y \subseteq V$, $X \cap Y = \emptyset$, we write $X \succeq_i Y$ if for each pair of agents $x \in X$ and $y \in Y$ we have $x \succeq_i y$.

A preference profile \mathcal{P} for V is a collection $(\succeq_i)_{i \in V}$ of preference lists for each agent $i \in V$. A profile \mathcal{P} may have the following properties:

1. It is *complete* if for each agent $i \in V$ it holds that $V_i \cup \{i\} = V$; otherwise it is *incomplete*.
2. The profile \mathcal{P} has *ties* if there is an agent $i \in V$ with ties in its preference list.

To an instance (V, \mathcal{P}) we assign an *acceptability graph* G , which has V as its vertex set and two agents are connected by an edge if each finds the other acceptable. Without loss of generality,

G does not contain isolated vertices, meaning that each agent has at least one agent which it finds acceptable.

Swaps and differences between two matchings. Given two preference lists \succeq and \succeq' , the *swap distance between \succeq and \succeq'* is defined as the number of pairs that are “ordered” differently; if \succeq and \succeq' are defined on different sets, then we assume that their swap distance is infinite. Formally,

$$\delta(\succeq, \succeq') := \begin{cases} \infty, & \succeq \text{ and } \succeq' \text{ are defined on different sets,} \\ |\{(x, y) \mid x \succ y \wedge y \succeq' x\}| + \\ |\{(x, y) \mid x \sim y \wedge (x, y) \notin \sim'\}|, & \text{otherwise.} \end{cases}$$

For example, if an agent has a preference list on $\{a, b, c\}$ where all three agents are tied on the first position, that is, $a \sim b \sim c$, then moving to the list $c \succ \{a, b\}$ requires two swaps.

Let \mathcal{P}_1 and \mathcal{P}_2 be two preference profiles for the same set V of agents. The *swap distance between \mathcal{P}_1 and \mathcal{P}_2* , denoted as $|\mathcal{P}_1 \oplus \mathcal{P}_2|$, is defined as the sum of the swap distances between the preference lists of each agent in the two preference profiles. Formally, let \succeq_i^j be the preference list of an agent $i \in V$ in the profile \mathcal{P}_j for $j = 1, 2$. We have that $|\mathcal{P}_1 \oplus \mathcal{P}_2| = \sum_{i \in V} \delta(\succeq_i^1, \succeq_i^2)$.

Given a set V of agents, a *matching* M of V is a set of pairwise disjoint pairs of agents in V . Given two matchings M_1 and M_2 for the same set V with $2n$ agents, we define the *distance between M_1 and M_2* as the size of the symmetric difference of M_1 and M_2 , formally: $\text{dist}(M_1, M_2) := |M_1 \Delta M_2| = |M_1 \setminus M_2| + |M_2 \setminus M_1|$.

Blocking pairs and stable matchings. Let a preference profile \mathcal{P} for a set V of agents, the corresponding acceptability graph G , and a matching $M \subseteq E(G)$ be given. For a pair $\{x, y\}$ of agents, if $\{x, y\} \in M$, then we denote the corresponding partner y by $M(x)$; otherwise we call this pair *unmatched*. We write $M(x) = \perp$ if agent x has *no partner*, i.e., if the agent x is not involved in any pair in M . We use $\perp(M)$ to denote the set of unmatched agents in a matching M , that is, $\perp(M) = \{x \mid M(x) = \perp\}$. If no agent x has $M(x) = \perp$ (i.e., $\perp(M) = \emptyset$), then M is called *perfect*.

Given a matching M of \mathcal{P} , an unmatched pair $\{x, y\} \in E(G) \setminus M$ is *blocking* M if both x and y prefer each other to being unmatched or to their assigned partners, i.e. it holds that $(M(x) = \perp \vee y \succ_x M(x)) \wedge (M(y) = \perp \vee x \succ_y M(y))$. We call a matching M *stable*³ if no unmatched pair is blocking M . The STABLE ROOMMATES problem is defined as follows:

STABLE ROOMMATES (SR)

Input: A preference profile \mathcal{P} for a set $V = \{1, 2, \dots, 2n\}$ of $2n$ agents.

Question: Does \mathcal{P} admit a stable matching?

STABLE ROOMMATES and STABLE MARRIAGE. The bipartite variant of STABLE ROOMMATES, called STABLE MARRIAGE, has as input two n -element disjoint sets $U \uplus W$ of agents such that each agent $u \in U$ from U has a *preference list on $W_u \subseteq W$* and each agent $w \in W$ from W has a *preference list on $U_w \subseteq U$* . By definition, the underlying acceptability graph of a STABLE MARRIAGE instance is bipartite. Accordingly, this instance has complete preferences if this graph is a complete bipartite graph.

The following fundamental result from the literature guarantees that we can deal with incomplete preferences without ties similarly to the case with complete preferences with ties.

Proposition 1 ([19, Theorem 1.4.2, Theorem 4.5.2]). *For incomplete preferences without ties, the whole agent set can be partitioned into two disjoint subsets R and S such that every stable matching matches every agent from R and none of the agents from S . For agent sets of size $2n$, this partition can be computed in $O(n^2)$ time.*

³We exclusively focus on the weak stability concept [28]. We conjecture that several results will also hold at least for strong stability. The proofs, however, may need non-trivial adjustments.

3 Algorithms for Incremental Stable Marriage and Roommates

As a warm-up, we first show that our most restricted problem variant, INCREMENTAL STABLE MARRIAGE without ties, can be solved in polynomial time. The main idea behind this result is based on the fact that there exists a compact and polynomial-time computable representation of all stable matchings, the so-called partially ordered set of polynomially many rotations [19, Chapter 2]. Herein, a rotation involves a subset of some stable matching “reducing” which results in another stable matching [19, Chapter 2.5.1]. Equipping these rotation with some appropriate weight, we can reduce our problem to finding a closed subset of rotations with maximum weight, which can be solved in polynomial-time using an approach similar to the one discussed in [19, Chapter 3.6.1]. We refer to Appendix A for proof details.

Proposition 2. INCREMENTAL STABLE MARRIAGE *without ties* can be solved in $O(n^3)$ time.

Proof. The result from Proposition 2 follows from Lemma 9 and Lemma 10. \square

Second we show that our least restricted NP-hard problem variant, INCREMENTAL STABLE ROOMMATES with ties allowed, can at least be solved in polynomial time when the distance between the matchings M_1 and M_2 is a constant. In other words, Proposition 3 presents an XP-algorithm for the distance as a parameter.

Proposition 3. INCREMENTAL STABLE ROOMMATES *with ties* can be solved in $n^{O(k)}$ time, where $k = |M_1 \Delta M_2|$.

Proof. Let $\hat{n} = |M_1|$. The algorithm first guesses positive integers k_1, k_2 with $k_1 + k_2 \leq k$ which we further treat as follows. We denote by k_1 the number of matching edges leaving M_1 (that is, $|M_1 \setminus M_2|$) and by k_2 the number of new edges (that is, $|M_2 \setminus M_1|$). Now, for each such pair (k_1, k_2) we first guess k_1 edges to delete from M_1 . Notice that there are $O(\hat{n}^{k_1}) = O(n^k)$ possible guesses. Let \hat{M}_1 be the rest of the matching M_1 (after we delete the just guessed edges). Then we guess k_2 pairs of vertices not matched in \hat{M}_1 . This completely defines the matching M_2 which can be checked in polynomial-time for stability. Notice that there are $\binom{2n-2\hat{n}+2k_1}{2k_2} \cdot \frac{(2k_2)!}{2 \cdot k_2!} = O(2k^{2k} \cdot (2n)^{2k})$ possible guesses. Repeating the above procedure for every pair (k_1, k_2) and applying the obvious bound $k \leq n$, we get the overall running time $n^{O(k)}$ as claimed. \square

We remark that Proposition 3 can also be shown by using an idea of Our approach, however, is much simpler. The most pressing question following from Proposition 3 is to ask for which cases this result can be improved to a fixed-parameter tractability result for the parameter k , the distance between M_1 and M_2 . Unfortunately, we will show in the next section that this is not possible when ties are involved. In the remainder of this section, however, we show that this is possible when ties are not allowed. Formally, we show the following.

Theorem 1. INCREMENTAL STABLE ROOMMATES *without ties* can be solved in $O(2^k n^4)$ time.

The algorithm behind Theorem 1 is partially inspired by the polynomial-time algorithm for solving MAXIMUM WEIGHT STABLE MARRIAGE [19, Chapter 2.5.1]. (also see Appendix A for more details). The high-level structure of our algorithm is as follows:

1. Using the algorithm of Irving [21] and based on the structural insights of Gusfield [18] that the set of all stable matchings can be compactly and efficiently represented via the so-called partially ordered set (poset) of polynomially many rotations, we compute the poset of rotations for the new preference profile P_2 (see Section 3.1).

Roughly speaking, a rotation (relative to a preference profile) involves a specific cyclic sequence σ of agents where the first acceptable partner in the preference list of each agent x in σ is exactly the second acceptable partner in the preference list of x 's predecessor

in σ . We remark that for a rotation, removing the first acceptable partner of each agent σ will not alter the existence of a stable matching.

2. To capture the (different) costs of eliminating one rotation from each dual pair (see [Definition 3](#)) with respect to the resulting distance between the matching M_1 and M_2 , we compute two weights for each rotation (see [Section 3.2](#)).
3. Finally, to consistently choose one rotation from each dual pair, and additionally respecting the weight constraints, we reduce our problem to an auxiliary problem to be defined below and provide a fixed-parameter algorithm for this auxiliary problem (see [Section 3.3](#)).

Each high-level step of the algorithm will be described in one of the subsequent subsections.

To define our auxiliary problem (which may be of independent interest), for each partially ordered set (R, \succeq) , we say that a given subset $C \subseteq R$ is *closed with respect to* the relation \succeq if for each pair $\{\rho, \sigma\} \subseteq R$ of elements it holds that $\rho \in C$ and $\sigma \succeq \rho$ imply $\sigma \in C$.

WEIGHTED CONFLICT-FREE CLOSED SUBSET

Input: A partially ordered set (R, \succeq) , an undirected graph $G = (R, E)$, a weight function $w: R \rightarrow \mathbb{N}$, a positive integer ℓ , and a budget $b \in \mathbb{N}$.

Question: Is there a closed subset $C \subseteq R$ of size ℓ which is independent in G such that $\sum_{c \in C} w(c) \leq b$?

Here, the conflicts are modeled by the edges so that the graph is also referred to as a *conflict graph* and seeking for an independent set solution means seeking for a *conflict-free set*. The structure of the rotation poset implies that in our case the graph G in fact consists of ℓ edges forming a matching of size ℓ . In particular, we will show that we can focus on the special case in which the conflictgraph G consists of ℓ disjoint cliques. For this case we give an algorithm for WEIGHTED CONFLICT-FREE CLOSED SUBSET for parameter $b + \Delta(G)$, where $\Delta(G)$ is the maximum vertex degree in the graph G . Moreover, it follows from our reduction procedure that $b \leq k$ (see [Section 3.3](#)).

3.1 Preprocessing and identification of the rotations

We first recall Irving's polynomial-time algorithm [\[21\]](#) for determining whether an instance of STABLE ROOMMATES without ties has a stable matching and fundamental structural properties behind all stable matchings [\[18\]](#). Irving's algorithm is divided into two phases: Phase 1 and Phase 2. Phase 1 involves a sequence of proposals from each agent i to the first agent j on i 's list, and each such proposal resulting in the deletion of all successors of i from j 's list. Phase 1 does not alter any stable matching since in this phase j is removed from i 's preference list (and i is removed from j 's preference list) only when $\{i, j\}$ does not form a pair in any stable matching. The set of lists at the end of the first phase is called *Phase 1 table* (please refer to [Example 1](#) for an illustration).

There are three possibilities for the Phase 1 table; note that the Phase 1 table is unique.

Every list in the table is empty. Then, there is no stable matching; note that Irving's original algorithm assumes that the input preferences are complete, implying that whenever a list becomes empty, then there is not stable matching; for incomplete preferences we can only infer the non-existence of any stable matching if every list becomes empty.

Every non-empty list contains exactly one agent. Then, we find a unique stable matching.

At least one list contains more than one agent. Then, we proceed to Phase 2.

For agents with empty preference lists in the Phase 1 table, we use the following.

Proposition 4 ([19, Theorem 4.5.2]). *For each agent, if its preference list becomes empty after Phase 1, then it will not be matched by any stable matching; otherwise it must be matched by all stable matchings.*

Since in the first two cases the instance of INCREMENTAL STABLE ROOMMATES is trivial to solve, we assume the third case in the following and we can ignore every agent whose list becomes empty after Phase 1. Notice that in the Phase 1 table we always have that if an agent i is ranked first in j 's preference list, then j is ranked the last in i 's preference list. This invariant we keep throughout the whole Phase 2. We need some definitions for Phase 2. A *preference table* is the Phase 1 table or any Phase 2 table (i.e., it is a collection of preference lists).

Definition 1 (Rotations). *A rotation exposed in a preference table \mathcal{T} is an ordered sequence*

$$(e_0, h_0), (e_1, h_1), \dots, (e_{r-1}, h_{r-1})$$

of pairs of agents such that h_i and $h_{i+1 \bmod r}$ are the first and the second agent on e_i 's list in \mathcal{T} , for all $0 \leq i \leq r-1$.

Note that if (e, h) is a pair in a rotation, then e is ranked last by h with respect to \mathcal{T} .

Definition 2 (Elimination of an exposed rotation). *The elimination of an exposed rotation $(e_0, h_0), (e_1, h_1), \dots, (e_{r-1}, h_{r-1})$ from table \mathcal{T} is the following operation. For every $0 \leq i \leq r-1$, remove every entry below e_i in h_{i+1} 's list in \mathcal{T} , i.e., move the bottom of h_{i+1} 's list up to e_i (from e_{i+1}). Then for each agent p who was just removed from h_{i+1} 's preference list, remove h_{i+1} from p 's list.*

In Phase 2, exposed rotations (see Example 1) are eliminated from the preference table one by one until some list becomes empty, which means there is no stable matching, or no rotation is exposed in the table, which means the list of every agent contains exactly one agent and we get a stable matching.

Proposition 5 ([22, Corollary 3.2]). *If some agent obtains an empty preference list in Phase 2, then there is no stable matching.*

To get better acquainted with Irving's algorithm and, more importantly, the notion of rotations, their exposition, and their elimination, we give an example illustrating this.

Example 1 ([18]). *Consider the following profile with eight agents—the complete profile is to the left and the Phase 1 table is to the right. For the sake of readability, the symbol \succ between the agents in the preference lists of each agent are omitted.*

1	7	2	6	8	5	3	4	1	2	6	5	3	4		
2	4	6	5	3	8	1	7	2	6	5	3	8	1		
3	5	2	1	7	4	6	8	3	5	2	1	7	4	6	
4	1	7	3	6	5	8	2	4	1	7	3	6	5	8	
5	7	1	8	4	6	2	3	5	7	1	8	4	6	2	3
6	7	3	8	4	5	1	2	6	3	8	4	5	1	2	
7	2	8	4	3	5	6	1	7	8	4	3	5			
8	4	2	3	5	6	7	1	8	4	2	5	6	7		

In the Phase 1 table, there are two rotations: $\rho_1 = ((1, 2), (2, 6), (3, 5))$ and $\rho_2 = ((4, 1), (5, 7))$. Now, we eliminate the rotation ρ_1 , that is, we remove

- the top choices for agents 1, 2, 3 and
- 8, 1 from the preference list of agent 2, since 2 finds 3 better than these two and is becoming the top choice for 3.

After the elimination of ρ_1 , we obtain the following table.

1	6	5	3	4		
2	5	3				
3	2	1	7	4	6	
4	1	7	3	6	5	8
5	7	1	8	4	6	2
6	3	8	4	5	1	
7	8	4	3	5		
8	4	5	6	7		

Now ρ_2 is still exposed, and there is a new exposed rotation $\rho_3 = ((2, 5), (6, 3), (7, 8), (8, 4))$.

If we continue to eliminate ρ_2 , then we obtain the following table.

1	6	5				
2	5	3				
3	2	4	6			
4	7	3	6	5	8	
5	1	8	4	6	2	
6	3	8	4	5	1	
7	8	4				
8	4	5	6	7		

Now ρ_3 is the only exposed rotation. We eliminate ρ_3 and obtain the following table.

1	6	5				
2	3					
3	2					
4	7					
5	1	8				
6	8	1				
7	4					
8	5	6				

Now $\rho_4 = ((1, 6), (8, 5))$ and $\rho_5 = ((5, 1), (6, 8))$ are exposed. If we continue to eliminate ρ_4 , we obtain the following table, where every person has exactly one entry on his list, and we get a stable matching.

1	5					
2	3					
3	2					
4	7					
5	1					
6	8					
7	4					
8	6					

There are two types of rotations, which can be used to characterize all stable matchings. Let D denote the execution tree when Phase 2 is executed in all possible ways on the Phase 1 table. Note that each node x in D represents a table $\mathcal{T}(x)$, which is the current table state of the algorithm at node x . Let R be the set of all rotations which are exposed in some table of some node in D .

Definition 3 (Dual and singleton rotations). If $\rho \in R$ with $\rho = ((e_0, h_0), (e_1, h_1), \dots, (e_{r-1}, h_{r-1}))$, then the negation of ρ is defined as $\neg\rho = ((h_0, e_{r-1}), (h_1, e_0), \dots, (h_{r-1}, e_{r-2}))$.

If $\neg\rho \in R$, then we call ρ and $\neg\rho$ a dual pair of rotations. Any rotation without a dual is called a singleton rotation.⁴

⁴The singleton rotation is called *singular rotation* in Gusfield and Irving [19], Manlove [28]; however, since we closely follow Gusfield [18], we prefer to use the term singleton rotation. This is mainly to make it easier for the reader when referring to lemmata of Gusfield.

Definition 4 (Partially ordered set, and closed and complete subest of rotations). *A subset $C \subseteq R$ of rotations is complete if it contains*

- (i) *all singleton rotations and*
- (ii) *exactly one rotation from each dual pair of rotation.*

A binary relation \triangleright on R is defined as follows. For each pair $\{\rho_1, \rho_2\}$ of rotations, if the elimination of ρ_1 is necessary for ρ_2 to be exposed (i.e., the elimination of ρ_1 precedes the exposition of ρ_2 on every path in D), then we say that ρ_1 precedes ρ_2 , written as $\rho_1 \triangleright \rho_2$. The partial order \succeq is the reflexive closure of the precedence relation \triangleright . A subset C of the rotation R is closed with respect to the partial order \succeq (in short, closed) if for each pair $\{\rho_1, \rho_2\}$ of rotations it holds that if $\rho_2 \in C$ and $\rho_1 \succeq \rho_2$, then $\rho_1 \in C$.

Note that the rotation poset \succeq on R is an $O(n^2)$ -sized representation of all stable matchings. It follows from Gusfield [18] that one can compute R in $O(n^3 \cdot \log(n))$ time.

Lemma 1 ([18]). *For a given STABLE ROOMMATES instance without ties, there is a one-to-one correspondence between the set of stable matchings and complete and closed subsets of the rotation poset (R, \succeq) . Finding this poset (R, \succeq) can be done in $O(n^3 \cdot \log(n))$ time, where n denotes the number of agents in the instance.*

According to Lemma 1, to solve our problem, it suffices to search for a complete and closed subset of rotations for the second input profile \mathcal{P}_2 such that the corresponding stable matching is closest to M_1 .

3.2 Proposal sets and rotation weights

In this section, we study the influence of a rotation elimination on the distance between the ultimate resulting matching and the initial matching. In order to do so, we first define proposal sets for preference tables and weights of rotations. Here, the weight of a rotation shall capture how many pairs of the initial matching we can additionally obtain if we eliminate this rotation. Then we study the properties of the thus introduced rotation weights. This is used later to finish the sought reduction and to determine upper bounds on parameter values of the resulting instance.

Imagine that in a preference table, every agent proposes to the first agent in their current preference list. Then, we get a set of ordered pairs, where the agent indicated by the first component of each pair proposes to the agent indicated by the second component in the pair.

Definition 5 (Proposal sets). *For each preference table \mathcal{T} , the proposal set for \mathcal{T} is defined as*

$$\mathcal{S}_{\mathcal{T}} := \{(i, j) \mid i \in U \text{ and } j \text{ is the first agent on } i\text{'s list in } \mathcal{T}\},$$

where (i, j) represents a proposal pair $i \rightarrow j$ (i.e., agent i proposes to agent j).

For each matching M , the proposal set \mathcal{S}_M for M is defined as $\mathcal{S}_M := \{(i, j), (j, i) \mid \{i, j\} \in M\}$.

By the definition of proposal sets, for each matching M we have

$$\text{dist}(M_1, M) = |M_1 \Delta M| = |M_1| + |M| - 2|M_1 \cap M| = |M_1| + |M| - |\mathcal{S}_{M_1} \cap \mathcal{S}_M|. \quad (1)$$

Hence, we are looking for a matching M_2 which is stable with respect to profile \mathcal{P}_2 such that

$$|\mathcal{S}_{M_1} \cap \mathcal{S}_{M_2}| \geq |M_1| + |M_2| - k. \quad (2)$$

Since every complete and closed rotation subset contains all singleton rotations and since all singleton rotations can be eliminated from the Phase 1 table before all dual rotations [18],

we can first eliminate all singleton rotations. By [Proposition 5](#), if after eliminating all singleton rotations, some agent obtains an empty preference list, then we can immediately conclude that the given profile does not admit any stable matching. Thus, in the following, we mainly work with $R_2 \subseteq R$ which is the set of all dual rotations.

To measure the benefit of eliminating a rotation, we define the following.

Definition 6 (Gain and lost proposal pairs after a rotation elimination). *For each rotation $\rho \in R$ with $\rho = (e_0, h_0), (e_1, h_1), \dots, (e_{r-1}, h_{r-1})$, the set of proposal pairs gained (lost) by eliminating rotation ρ is defined as follows.*

$$S_\rho^+ := \{(e_0, h_1), (e_1, h_2), \dots, (e_{r-1}, h_0)\},$$

$$S_\rho^- := \{(e_0, h_0), (e_1, h_1), \dots, (e_{r-1}, h_{r-1})\}.$$

Further, let $w^+(\rho) := |S_\rho^+ \cap S_{M_1}|$ and $w^-(\rho) := |S_\rho^- \cap S_{M_1}|$ be the number of proposal pairs gained and lost by the elimination of rotation ρ .

Observe that the two sets S_ρ^+ and S_ρ^- are independent of the table \mathcal{T} and all of the proposal pairs of agents not involved in the rotation ρ remain the same before and after the elimination of ρ .

Since in order to obtain a stable matching, we have to eliminate exactly one rotation from each dual pair of rotations, we now prove that their weights are complementary.

Lemma 2. *Let $\rho \in R$ be a dual rotation, then $w^+(\rho) = w^-(\neg\rho)$.*

Proof. Let $\rho := (e_0, h_0), (e_1, h_1), \dots, (e_{r-1}, h_{r-1})$, and let S_ρ^+ and S_ρ^- be the sets of proposal pairs gained and lost by eliminating ρ , respectively. That is,

$$S_\rho^+ = \{(e_0, h_1), (e_1, h_2), \dots, (e_{r-1}, h_0)\}, \text{ and} \quad (3)$$

$$S_\rho^- = \{(e_0, h_0), (e_1, h_1), \dots, (e_{r-1}, h_{r-1})\}. \quad (4)$$

By the definition of dual rotations, the dual of ρ is equal to $\neg\rho = (h_0, e_{r-1}), (h_1, e_0), \dots, (h_{r-1}, e_{r-2})$. Accordingly, we have:

$$S_{\neg\rho}^+ = \{(h_0, e_0), (h_1, e_1), \dots, (h_{r-1}, e_{r-1})\}, \text{ and} \quad (5)$$

$$S_{\neg\rho}^- = \{(h_0, e_{r-1}), (h_1, e_0), \dots, (h_{r-1}, e_{r-2})\}. \quad (6)$$

By (3) and (6), we can infer that

$$\text{for each ordered pair } (i, j) \text{ of agents it holds that } (i, j) \in S_\rho^+ \text{ if and only if } (j, i) \in S_{\neg\rho}^-. \quad (7)$$

Since M_1 is a matching, it follows that for each unordered pair $\{i, j\} \in M_1$ of agents we have that $(i, j), (j, i) \in S_{M_1}$. Together with (7), we conclude that

$$\text{for each ordered pair } (i, j) \text{ of agents it holds that } (i, j) \in S_\rho^+ \cap S_{M_1} \text{ iff. } (j, i) \in S_{\neg\rho}^- \cap S_{M_1}.$$

Thus,

$$w^+(\rho) = |S_\rho^+ \cap S_{M_1}| = |S_{\neg\rho}^- \cap S_{M_1}| = w^-(\neg\rho).$$

□

3.3 Reduction to Weighted Conflict-Free Closed Subset

In this subsection we give a reduction from INCREMENTAL STABLE ROOMMATES to WEIGHTED CONFLICT-FREE CLOSED SUBSET in which the conflict graph is a union of disjoint edges. In order to do so, we show that the distance between the target (i.e., initial) matching M_1 and a matching M_C resulting from the elimination of a complete and closed set C of rotations for \mathcal{P}_2 (if such set exists) depends only on $\sum_{\rho \in C} w^-(\rho)$.

In the remainder of the section, let S_0 be the proposal set for the table obtained from the Phase 1 table for \mathcal{P}_2 followed by elimination of all singleton rotations to it and let R_2 be the set of all dual rotations for \mathcal{P}_2 ; it is known that R_2 has at most $\binom{n}{2}$ rotations.

Lemma 3 ([18, Corollary 5.1]). $|R_2| \leq \binom{n}{2}$.

Upper-bounding the sum of weights of a complete and closed subset. Before we continue with the procedure, we compare the sizes of M_C and M_1 ; recall that M_C is a matching resulting from eliminating a complete and closed set $C \subseteq R$. By Proposition 4, every agent that has non-empty list after Phase 1 must be matched under M_C and these agents are exactly those agents who hold some proposals in S_0 . Consequently, for each agent x that is matched under M_1 but does not hold a proposal under S_0 it holds that $\{x, M_1(x)\} \in M_1 \Delta M_C$. Thus, we define the following reduction rule.

Reduction Rule 1. For each agent x matched under M_1 , if for each agent y it holds that $(x, y) \notin S_0$, then delete $\{x, M_1(x)\}$ from M_1 and decrease k by one.

Lemma 4. *Rule 1 is sound and can be implemented in $O(n^2)$ time. Moreover, if Rule 1 does not apply, then for each stable matching M of profile \mathcal{P}_2 it holds that $|M_1| \leq |M| = |S_0|/2$.*

Proof. Let x be as defined in the rule, i.e., x is matched under M_1 but does not hold a proposal in S_0 . For ease of notation, let $M_1^- := M_1 \setminus \{\{x, M_1(x)\}\}$. Let M be a stable matching for \mathcal{P}_2 . By Proposition 4, it follows that x is unmatched under M . Thus, $M_1 \Delta M = (M \Delta M_1^-) \cup \{x, M_1(x)\}$. This implies that $|M_1 \Delta M| \leq k$ if and only if $|M_1^- \Delta M| \leq k - 1$. The soundness of the reduction rule follows.

As for the running time, for $2n$ agents in \mathcal{P}_2 , in $O(n^2)$ time we can complete Phase 1 and obtain the proposal set S_0 which has size at most $2n$. Then, in $O(n)$ time we can check for each matched agent (under M_1) whether it is “unmatched” in S_0 .

As for the second statement, it is immediate that $|M_1| \leq |M|$. Again, by Proposition 4, it holds that $|M| = |S_0|/2$. Hence, $|M_1| \leq |M| = |S_0|/2$. \square

Next, we upper-bound the size of the intersection between the target (i.e., initial) matching and the sought matching.

Lemma 5. *Let C be a complete and closed subset of rotations in R_2 and let M_C be the stable matching associated with C . Then, the following holds.*

- (i) $|S_{M_1} \cap S_{M_C}| = |S_{M_1} \cap S_0| + \sum_{\sigma \in R_2} w^+(\sigma) - 2 \cdot \sum_{\rho \in C} w^-(\rho)$.
- (ii) $\text{dist}(M_1, M_C) \leq k$ if and only if $\sum_{\rho \in C} w^-(\rho) \leq \frac{|S_{M_1} \cap S_0| + \sum_{\sigma \in R_2} w^+(\sigma) - |M_1| - |M_C| + k}{2}$.
- (iii) $|S_{M_1} \cap S_0| + \sum_{\sigma \in R_2} w^+(\sigma) - |M_1| - |M_C| \leq 0$.

Proof. To show the first statement, let us first observe the following, which is similar to the case in the marriage setting (see Lemma 8 and Corollary 1).

Claim 1. *Let \mathcal{T} be a preference table and ρ be a rotation exposed in \mathcal{T} , and let \mathcal{T}' be the preference table obtained from \mathcal{T} by eliminating ρ . Then, $|S_{M_1} \cap S_{\mathcal{T}'}| = |S_{M_1} \cap S_{\mathcal{T}}| + w^+(\rho) - w^-(\rho)$.*

Proof. The proof is similar to the one given for Lemma 8. Let $\rho := ((e_0, h_0), \dots, (e_{r-1}, h_{r-1}))$ be a rotation exposed in the table \mathcal{T} . In the following, all subscripts $i + 1$ are taken modulo r . By the definition of \mathcal{T}' and the definition of proposal sets we have that

$$\begin{aligned} \mathcal{S}_{\mathcal{T}'} &= \{(x, y) \in \mathcal{S}_{\mathcal{T}} \mid (x, y) \notin \rho\} \uplus \{(e_i, h_{i+1}) \mid 0 \leq i \leq r-1\} \\ &= (\mathcal{S}_{\mathcal{T}} \cup \mathcal{S}_{\rho}^+) \setminus \mathcal{S}_{\rho}^-. \end{aligned} \quad (8)$$

Thus, we prove the statement in the claim by showing the following.

$$\begin{aligned} |\mathcal{S}_{M_1} \cap \mathcal{S}_{\mathcal{T}'}| &\stackrel{(8)}{=} |\mathcal{S}_{M_1} \cap ((\mathcal{S}_{\mathcal{T}} \cup \mathcal{S}_{\rho}^+) \setminus \mathcal{S}_{\rho}^-)| \\ &= |((\mathcal{S}_{M_1} \cap \mathcal{S}_{\mathcal{T}}) \cup (\mathcal{S}_{M_1} \cap \mathcal{S}_{\rho}^+)) \setminus (\mathcal{S}_{M_1} \cap \mathcal{S}_{\rho}^-)| \\ &= |(\mathcal{S}_{M_1} \cap \mathcal{S}_{\mathcal{T}})| + w_{\rho}^+ - w_{\rho}^-. \end{aligned}$$

The last equation holds because $\mathcal{S}_{\mathcal{T}} \cap \mathcal{S}_{\rho}^+ = \emptyset$ and $\mathcal{S}_{\rho}^- \subseteq \mathcal{S}_{\mathcal{T}}$ and because of Definition 6. (of Claim 1) \diamond

By applying the above repeatedly, we obtain the following for an (arbitrary) closed subset of rotations.

Claim 2. *If C' is a closed subset of rotations and \mathcal{T} is the preference table obtained by eliminating all rotations from C' on the Phase 1 table, then $|\mathcal{S}_{M_1} \cap \mathcal{S}_{\mathcal{T}}| = |\mathcal{S}_{M_1} \cap \mathcal{S}_0| + \sum_{\rho \in C'} w^+(\rho) - \sum_{\rho \in C'} w^-(\rho)$.*

Now, we are ready to show our first statement. To this end, let C and M_C be as defined in the lemma. By Claim 2, it follows that

$$\begin{aligned} |\mathcal{S}_{M_1} \cap \mathcal{S}_{M_C}| &= |\mathcal{S}_{M_1} \cap \mathcal{S}_0| + \sum_{\rho \in C} w^+(\rho) - \sum_{\rho \in C} w^-(\rho) \\ &= |\mathcal{S}_{M_1} \cap \mathcal{S}_0| + \left(\sum_{\sigma \in R_2} w^+(\sigma) - \sum_{\sigma \in R_2 \setminus C} w^+(\sigma) \right) - \sum_{\rho \in C} w^-(\rho) \\ &\stackrel{\text{Lemma 2}}{=} |\mathcal{S}_{M_1} \cap \mathcal{S}_0| + \left(\sum_{\sigma \in R_2} w^+(\sigma) - \sum_{\sigma \in R_2 \setminus C} w^-(\neg\sigma) \right) - \sum_{\rho \in C} w^-(\rho) \\ &\stackrel{C \text{ is complete}}{=} |\mathcal{S}_{M_1} \cap \mathcal{S}_0| + \sum_{\sigma \in R_2} w^+(\rho) - 2 \cdot \sum_{\rho \in C} w^-(\rho). \end{aligned}$$

This completes the proof for the first statement.

The second statement follows directly from (1), (2), and the first statement.

As for the last statement, we first observe that for each two distinct rotations σ and σ' from R_2 it holds that $\mathcal{S}_{\sigma}^+ \cap \mathcal{S}_{\sigma'}^+ = \emptyset$. To see this, let $\sigma := ((e_0, h_0), \dots, (e_{r-1}, h_{r-1}))$ which is exposed in some preference table \mathcal{T} . Then, by definition, it holds that $\mathcal{S}_{\sigma}^+ = \{(e_i, h_{i+1}) \mid 0 \leq i \leq r-1\}$. However, by [18, Lemma 5.1, Corollary 5.1], for each $i \in \{0, \dots, r-1\}$, no other other rotation ρ exists such that $(e_i, h_{i+1}) \in \mathcal{S}_{\rho}^+$.

Next, we observe that $\mathcal{S}_0 \cap \mathcal{S}_{\sigma}^+ = \emptyset$. To see this, let $\sigma := ((e_0, h_0), \dots, (e_{r-1}, h_{r-1}))$. Then, $\mathcal{S}_{\sigma}^+ = \{(e_i, h_{i+1}) \mid 0 \leq i \leq r-1\}$ ($i+1$ are taken modulo r). By the definition of rotations, it follows that e_i prefers h_i to h_{i+1} , implying that h_{i+1} is *not* the first agent in the preference list of e_i right after Phase 1. That is, $(e_i, h_{i+1}) \notin \mathcal{S}_0$. Thus, we have that

$$\begin{aligned} |\mathcal{S}_{M_1} \cap \mathcal{S}_0| + \sum_{\sigma \in R_2} w^+(\sigma) &= |\mathcal{S}_{M_1} \cap \mathcal{S}_0| + \sum_{\sigma \in R_2} |\mathcal{S}_{M_1} \cap \mathcal{S}_{\sigma}^+| = |\mathcal{S}_{M_1} \cap (\mathcal{S}_0 \cup \bigcup_{\sigma \in R_2} \mathcal{S}_{\sigma}^+)| \\ &\leq 2 \cdot |M_1|. \end{aligned} \quad (9)$$

Now, to show the last statement, it suffices to show that $|M_1| \leq |M_C|$. This is true by Lemma 4:

$$|\mathcal{S}_{M_1} \cap \mathcal{S}_0| + \sum_{\sigma \in R_2} w^+(\sigma) - |M_1| - |M_C| \stackrel{(9)}{\leq} 2 \cdot |M_1| - |M_1| - |M_C| \stackrel{\text{Lemma 4}}{\leq} 0. \quad \square$$




Figure 1: A diagram for the partial order \succeq on the rotation set R_2 for the profile given [Example 1](#), as discussed in [Example 2](#).

This allows us to reduce the given instance (in polynomial time) to an instance of WEIGHTED CONFLICT-FREE CLOSED SUBSET as follows; see [Example 2](#) for an illustration.

Construction 1. Finally, we arrive at the following instance of WEIGHTED CONFLICT-FREE CLOSED SUBSET:

- (i) Apply [Rule 1](#) in $O(n^2)$ time, and compute in $O(n^3 \cdot \log n)$ the rotation poset (R_2, \succeq) for \mathcal{P}_2 (see [Lemma 1](#)).
- (ii) Compute S_0 for the profile \mathcal{P}_2 . Let $R := R_2$ be the set of all dual rotations; note that $|R_2| \leq \binom{n}{2}$ ([Lemma 3](#)),
- (iii) Let G be a graph on R in which two elements of R are adjacent if they form a dual pair of rotations (consequently, G is a union of $\ell = |R|/2$ disjoint edges).
- (iv) The weight function w is defined by w^- .
- (v) The budget b on the sum of weights is $b := \frac{|S_{M_1 \cap S_0}| + \sum_{\sigma \in R_2} w^+(\sigma) - |M_1| - |S_0|/2 + k}{2}$.

Note that b is derived from [Lemma 5\(ii\)](#) such that we are searching for a complete and closed subset C of rotations whose sum of weights is bounded by b . To see this, since $|S_M| = |S_0|/2$ ([Lemma 4](#)), the budget b is in fact equal to $\frac{|S_{M_1 \cap S_0}| + \sum_{\sigma \in R_2} w^+(\sigma) - |M_1| - |M| + k}{2}$, where M is an arbitrary stable matching for \mathcal{P}_2 . Thus, by [Lemma 5\(ii\)](#), the budget b is at most $k/2$. Note also that the sum of the weights $w^-(\rho)$ is upper-bounded by n and thus the reduction presented above is a polynomial (many-to-one) reduction.

Example 2. Consider the profile given in [Example 1](#). Let $M_1 = \{\{1, 7\}, \{2, 3\}, \{4, 6\}, \{5, 8\}\}$ and $k = 4$. To find a stable matching M such that $\text{dist}(M_1, M) \leq 4$, we first compute the rotation poset $(R = \{\rho_1, \rho_2, \rho_3, \rho_4, \rho_5, \rho_6\}, \succeq)$. The partial order \succeq is shown in [Figure 1](#).

Since ρ_1 and ρ_3 are singleton rotations, we can first eliminate them and get the set of all dual rotations $R_2 = \{\rho_2, \rho_4, \rho_5, \rho_6\}$, where $\rho_2 = \neg \rho_6$ and $\rho_4 = \neg \rho_5$. After the elimination of ρ_1 and ρ_3 , we obtain the following table.

1	6	5	3	4
2	3			
3	2			
4	1	7		
5	7	1	8	
6	8	4	5	1
7	4	3	5	
8	5	6		

Now the proposal set $S_0 = \{(1, 6), (2, 3), (3, 2), (4, 1), (5, 7), (6, 8), (7, 4), (8, 5)\}$. Since $M_1 = \{\{1, 7\}, \{2, 3\}, \{4, 6\}, \{5, 8\}\}$, we have that $S_{M_1} = \{(1, 7), (7, 1), (2, 3), (3, 2), (4, 6), (6, 4), (5, 8), (8, 5)\}$. Then we can define weight functions w^+ and w^- . For example, for rotation $\rho_4 = ((1, 6), (8, 5))$:




Figure 2: The directed graph G on the rotation set R_2 for as discussed in [Example 2](#), which shows the precedence relation between the rotations. The weights of the weight function w are depicted next to the vertices (i.e., rotations).

$$\begin{aligned} S_{\rho_4}^+ &= \{(1, 5), (8, 6)\}, \text{ and} \\ S_{\rho_4}^- &= \{(1, 6), (8, 5)\}. \end{aligned}$$

Hence,

$$\begin{aligned} w^+(\rho_4) &= |S_{\rho_4}^+ \cap S_{M_1}| = |\{\emptyset\}| = 0, \text{ and} \\ w^-(\rho_4) &= |S_{\rho_4}^- \cap S_{M_1}| = |\{(8, 5)\}| = 1. \end{aligned}$$

Similarly, we can compute weights for other rotations in R_2 :

$$w^+(\rho_2) = 0, w^-(\rho_2) = 0, w^+(\rho_5) = 1, w^-(\rho_5) = 0, w^+(\rho_6) = 0, w^-(\rho_6) = 0.$$

Now we can construct the instance $(R = R_2, G = (R_2, E), w, \ell, b)$ of WEIGHTED CONFLICT-FREE CLOSED SUBSET. Edge set E consists of two edges $\{\rho_2, \rho_6\}$ and $\{\rho_4, \rho_5\}$ since they are dual pairs of rotations. The weight function w is defined as:

$$w(\rho_2) = w^-(\rho_2) = 0, w(\rho_4) = w^-(\rho_4) = 1, w(\rho_5) = w^-(\rho_5) = 0, w(\rho_6) = w^-(\rho_6) = 0.$$

Since $|R_2| = 4$, we have $\ell = |R_2|/2 = 2$. The budget b is given by:

$$\begin{aligned} b &= (|S_{M_1} \cap S_0| + \sum_{\sigma \in R_2} w^+(\sigma) - |M_1| - |S_0|/2 + k)/2 \\ &= (3 + 1 - 4 - 4 + 4)/2 \\ &= 0 \end{aligned}$$

Now the task is to find a closed subset $C \subseteq R$ of size $\ell = 2$ which is independent in G such that $\sum_{c \in C} w(c) \leq 0$. It is easy to see that $C = \{\rho_2, \rho_5\}$ is the only solution such that $\sum_{c \in C} w(c) \leq 0$. By eliminating ρ_2 and ρ_5 , we get matching $M_{\{\rho_1, \rho_2, \rho_3, \rho_5\}} = \{(1, 6), (2, 3), (7, 4), (8, 5)\}$. It is easy to check that $\text{dist}(M_1, M_{\{\rho_1, \rho_2, \rho_3, \rho_5\}}) = 4$.

Solving Weighted Conflict-Free Closed Subset when the conflict graph consists of ℓ disjoint cliques. Now, to prove [Theorem 1](#), we only need to show the following; recall that the budget b , as defined in the construction, is at most $k/2$.

Lemma 6. *If G consists of exactly ℓ cliques, then WEIGHTED CONFLICT-FREE CLOSED SUBSET can be solved in $O((\Delta(G) + 1)^b \cdot |R|^2)$ time.*

Proof. We present an algorithm for the special case of WEIGHTED CONFLICT-FREE CLOSED SUBSET when the conflict graph G is a union of exactly ℓ cliques (see [Algorithm 1](#)).

First we introduce some notation. For every element $p \in R$, let

$$T_p^\uparrow = \{q \in R \mid q \supseteq p\} \quad \text{and} \quad T_p^\downarrow = \{q \in R \mid p \supseteq q\}$$

be the set of all predecessors and the set of all ancestors of p (including p itself). For a set $P \subseteq R$ we define T_P^\uparrow as the union $\bigcup_{p \in P} T_p^\uparrow$; the set T_P^\downarrow is defined analogously.

Before we continue with the proof, we observe the following.

Observation 1. Let S be a solution to WEIGHTED CONFLICT-FREE CLOSED SUBSET. If $p \in S$, then $T_p^\uparrow \subseteq S$ and $T_{N_G(p)}^\downarrow \cap S = \emptyset$.

Proof. When $p \in S$, since S is closed, we get $T_p^\uparrow \subseteq S$. Now, due to the conflicts $S \cap N_G(p) = \emptyset$ and consequently no element of $T_{N_G(p)}^\downarrow$ can be a part of S . (of [Observation 1](#)) \diamond

According to [Observation 1](#), when we decide to include p into S , we have to also include all elements in T_p^\uparrow into S and delete all elements in T_p^\uparrow and $T_{N_G(p)}^\downarrow$ from R . We extend the weight of a single element to the set $T \subseteq R$ of elements in the natural way, that is, we set $w(T) := \sum_{p \in T} w(p)$.

To upper-bound the weights of the sets T_p^\uparrow from above, we present the following data reduction rules.

Reduction Rule 2. If $w(T_p^\uparrow) > b$, then we delete T_p^\downarrow from R .

Reduction Rule 3. If [Rule 2](#) is not applicable and there exists a clique with exactly one vertex p , then we add T_p^\uparrow to S and decrease b by its weight $w(T_p^\uparrow)$.

Reduction Rule 4. Let K be a clique in G . If $p \in T_K^\uparrow$, then we add T_p^\uparrow to S and decrease b by its weight $w(T_p^\uparrow)$.

In fact it follows from the results of Gusfield [18] that [Rule 4](#) is never used in any WEIGHTED CONFLICT-FREE CLOSED SUBSET instance resulting from the construction we just defined.

Claim 3. *Rules 2 to 4 are sound. Furthermore, Rules 2 to 4 can be implemented to run in $O(|R|^2)$ time.*

Proof. Note that we have to take at least one vertex from each clique in G into S since we are about to take ℓ elements in total; thus, [Rules 3](#) and [4](#) are sound. Clearly, no solution S can contain any set T of weight $w(T) > b$, yielding soundness of [Rule 2](#).

Indeed, [Rule 2](#) runs in $O(R)$ time. As for [Rules 3](#) and [4](#), we need to update T_q^\uparrow , T_q^\downarrow and the corresponding weights for every element q left in the (updated) set of rotations R . This can be done in a straightforward way in $O(|R|^2)$ time. (of [Claim 3](#)) \diamond

We apply [Rules 2](#) to [4](#) exhaustively. Now, we branch on a clique K with $0 < w(T_p^\uparrow) \leq b$ for every $p \in K$ since we have to add one element of K into any solution anyway. Note that in this branching we omit those cliques K for which $\min_{p \in K} w(T_p^\uparrow) = 0$ holds. This procedure yields a search tree of depth at most b with branching factor $\Delta(G) + 1$. Suppose that there is a leaf with nonnegative budget in which there is no clique left for branching. We claim that this is a yes-instance. At this point any clique left in G contains at least one element p with $w(T_p^\uparrow) = 0$; we select greedily these elements into the solution S . Observe that this gives a valid solution since all of these sets are of total weight 0 (note that by including p in the solution all of the weights can only decrease).

As for the claimed running time, recall that it is possible to update all of the sets T left for branching in $O(|R|^2)$ time. Furthermore, [Rules 3](#) and [4](#) are invoked $O(b)$ times and [Rule 2](#) at most once. \square

We finally come to our main theorem.

Algorithm 1:

```
1 FindS( $R, G, b, \ell$ )
2   if  $b < 0$  then return No;
3    $S \leftarrow \emptyset$ ;
4   while  $\exists p \in R$  with  $w(T_p^\uparrow) > b$  do delete  $T_p^\downarrow$  from  $R$  and  $G$ ; ▷ Rule 2
5   while  $\exists p \in R$  with  $p \in T_K^\uparrow$  for some clique  $K$  in  $G$  do ▷ Rule 4
6      $S \leftarrow S \cup T_p^\uparrow$ ;
7     delete  $T_p^\uparrow$  from  $R$  and  $G$ ;
8      $b \leftarrow b - w(T_p^\uparrow)$ ;
9   if  $G$  contains less than  $\ell$  cliques then return No;
10  while  $G$  contains an isolated vertex  $p$  do ▷ Rule 3
11     $S \leftarrow S \cup T_p^\uparrow$ ;
12    delete  $T_p^\uparrow$  from  $R$  and  $G$ ;
13     $b \leftarrow b - w(T_p^\uparrow)$ ;
14     $\ell \leftarrow \ell - 1$ ;
15  while  $\exists K$  clique in  $G$  with  $w(T_p^\uparrow) > 0$  for all  $p \in K$  do
16    foreach  $p$  in  $K$  do
17       $S' \leftarrow \text{FindS}(R \setminus (T_p^\uparrow \cup T_{K-p}^\downarrow), G, b - w(T_p^\uparrow), \ell - 1)$ ;
18      if  $S' \neq \text{No}$  and  $S \cup T_p^\uparrow \cup S'$  is independent in  $G$  then return  $S \cup T_p^\uparrow \cup S'$ ;
19    return No;
20  if  $\exists p \in R$  in clique  $K$  with  $w(T_p^\uparrow) = 0$  then return  $S \cup$ 
    FindS( $R \setminus (T_p^\uparrow \cup T_{K-p}^\downarrow), G, b, \ell - 1$ ); ▷  $R \neq \emptyset$ 
21  return  $S$ ;
```

Proof of Theorem 1. By [Construction 1](#), in polynomial time, we construct an instance for WEIGHTED CONFLICT-FREE CLOSED SUBSET where the budget b is upper-bounded by $k/2$ and the conflict graph consists of $|R_2|/2$ edges (see the reasoning right after [Construction 1](#)). By [Lemma 6](#), we can solve this instance and thus our problem in $O(2^k \cdot n^4)$ time. \square

4 Hard cases of Incremental Stable Roommates

Throughout this section, we are using the following non-standard “incremental” (resp. “decremental”) variants of the INDEPENDENT SET (resp. CLIQUE) problem to show parameterized intractability.

Our first problem asks for an independent set of size h for some graph in the case when an independent set of size h for the graph minus an edge is already known.

EDGE-INCREMENTAL INDEPENDENT SET

Input: A graph G , a distinguished edge $e^* \in E(G)$, a positive integer h , and an independent set S^* of size h for $G - e^*$.

Question: Does there exist an independent set S of size h in G ?

Our second problem asks for a clique with pendant edges of size h for some graph in the case when a clique of size h with pendant edges for the graph with an additional edge is known. A *clique with pendant edges for a graph G* is a subset $V' \subset V(G)$ of vertices such that V' forms a clique in G , i.e., each two vertices in V' are adjacent, and that each vertex in V' has at least one

neighbor outside V' . The size of a *clique with pendant edges* is defined as the number of vertices in the clique.

EDGE-DECREMENTAL CLIQUE WITH PENDANT EDGES

Input: A graph $G = (V, E)$, a distinguished edge $e^* \in E(G)$, a positive integer h , and a clique $S^* \subseteq V$ with pendant edges of size h for G .

Question: Is there a clique $S \subseteq V$ with pendant edges of size h in $G - e^*$?

Lemma 7. EDGE-INCREMENTAL INDEPENDENT SET *and* EDGE-DECREMENTAL CLIQUE WITH PENDANT EDGES *are NP-hard and W[1]-hard with respect to h .*

Proof. Note that W[1]-hardness under Turing-reductions is immediate for these problems and the simple tricks used to show the following lemma are to obtain hardness also under many-one reductions.

Edge-Incremental Independent Set. We give a reduction from INDEPENDENT SET parameterized by the size h of the solution to EDGE-INCREMENTAL INDEPENDENT SET again with parameter h . Let (\hat{G}, h) be an instance of INDEPENDENT SET. We construct graph G from $\hat{G} = (V, E)$ as follows: We add a set S^* of h new vertices and for each vertex $v \in V$ and each vertex $s \in S^*$ we add edge $\{v, s\}$. We finish the construction by picking two distinct vertices s and s' from S^* and adding to E the edge $e^* = \{s, s'\}$. Clearly, S^* is an independent set of size h in $G - e^*$ as required.

Observe that for each independent set S in G we either have $S \subseteq V$ or $S \subseteq S^*$, since we have added a complete bipartite graph between V and S . Now, since any independent set S in G with $S \subseteq S^*$ contains at most $h - 1$ vertices, an independent set of size h in G can only contain vertices from the set V , thus it must be an independent set of size h in the graph \hat{G} .

Edge-Decremental Clique with Pendant Edges. Without loss of generality, we assume that $h > 2$ and describe a straight-forward parameterized reduction from EDGE-INCREMENTAL INDEPENDENT SET parameterized by the size h of the solution to EDGE-DECREMENTAL CLIQUE WITH PENDANT EDGES with parameter h . Given the instance (G, e^*, S^*, h) of EDGE-INCREMENTAL INDEPENDENT SET, we create the graph G' by complementing the graph $G - e^*$ and adding for each original vertex in $v \in V(G')$ a new vertex that is only connected to v . It is easy to verify that (G', e^*, S^*, h) is a yes-instance of EDGE-DECREMENTAL CLIQUE WITH PENDANT EDGES if and only if (G, e^*, S^*, h) is a yes-instance of EDGE-INCREMENTAL INDEPENDENT SET. (The newly added vertices ensure the existence of the pendant edges and can never be part of a clique of size more than two.) \square

4.1 Incremental Stable Roommates is generally hard even without ties

To show that INCREMENTAL STABLE ROOMMATES without ties is NP-hard, we identify a relation of it to an egalitarian variant of stable matching where the egalitarian cost is minimized. Here, the *egalitarian cost* of a matching M is defined as the sum of the ranks of the agents with respect to their partners, and the *rank* of an agent x with respect to its partner $M(y)$ is equal to the number agents that x prefer over y . Feder [12] showed that finding a stable matching with minimum egalitarian cost is NP-hard for STABLE ROOMMATES, even for complete preferences without ties (see the work of Chen et al. [7] for fixed-parameter tractability results on this problem).

The original hardness proof by Feder [12] is to reduce from VERTEX COVER, which, given an undirected graph G and an integer $h' \in \mathbb{N}$, asks whether G admits a *vertex cover* of size h' , i.e., a size- h' vertex subset of $V' \subseteq V(G)$ such that each edge in $E(G)$ is incident to at least one vertex from V' . The basic idea behind the reduction is that putting a vertex to the solution set is equivalent to increasing the egalitarian cost by one. This correspondence can also be achieved in INCREMENTAL STABLE ROOMMATES by choosing an initial matching M_1 which is associated to an empty vertex set, thus finding a stable matching M_2 closest to M_1 is equivalent to finding

a vertex cover of minimum size. Note that VERTEX COVER and INDEPENDENT SET are dual to each other, i.e., a vertex subset V' is a vertex cover of size h' if and only if $V(G) \setminus V'$ is an independent set of size $|V(G)| - h'$. Due to this and since INDEPENDENT SET is $W[1]$ -hard with respect to $|V| - h'$, we will directly reduce from INDEPENDENT SET, also showing parameterized intractability for our problem.

Proposition 6. INCREMENTAL STABLE ROOMMATES *without ties* is NP-hard and $W[1]$ -hard with respect to $k' = |M_1 \cap M_2|$.

Proof Sketch. We present a reduction from the INDEPENDENT SET problem which is $W[1]$ -hard when parameterized by the solution size. Let $G = (V, E)$ be a graph and h be the desired size of the independent set.

Following essentially the construction of Feder [12], we create our instance of INCREMENTAL STABLE ROOMMATES as follows. For each vertex $v_i \in V$, we create four agents $p_i, \bar{p}_i, q_i, \bar{q}_i$. The preferences of these agents in \mathcal{P}_2 are identical to the profile constructed by Feder [12]:

$$\begin{aligned} \text{agent } p_i: \bar{p}_i \succ [\{p_j \mid \{p_i, p_j\} \in E\}] \succ \bar{q}_i, & \quad \text{agent } \bar{p}_i: q_i \succ p_i, \\ \text{agent } q_i: \bar{q}_i \succ \bar{p}_i, & \quad \text{agent } \bar{q}_i: p_i \succ q_i. \end{aligned}$$

The old profile \mathcal{P}_1 is defined as follows.

$$\begin{aligned} \text{agent } p_i: \bar{p}_i \succ \bar{q}_i \succ [\{p_j \mid \{p_i, p_j\} \in E\}], & \quad \text{agent } \bar{p}_i: q_i \succ p_i, \\ \text{agent } q_i: \bar{q}_i \succ \bar{p}_i, & \quad \text{agent } \bar{q}_i: p_i \succ q_i. \end{aligned}$$

Following the proof of Feder [12] we can conclude that matching all p_i with the \bar{q}_i and q_i with the \bar{p}_i is a stable matching for \mathcal{P}_1 . In order to obtain a stable matching for \mathcal{P}_2 , however, one can only keep those p_i matched with the \bar{q}_i that correspond to an independent set. \square

4.2 Ties in the preferences make Incremental Stable Marriage hard

We show that INCREMENTAL STABLE MARRIAGE becomes intractable when ties are allowed even if the two preference profiles \mathcal{P}_1 and \mathcal{P}_2 are almost identical. The following result is achieved by a parameterized reduction from EDGE-DECREMENTAL CLIQUE WITH PENDANT EDGES.

Theorem 2. INCREMENTAL STABLE MARRIAGE *with ties* is $W[1]$ -hard with respect to k , even if $|\mathcal{P}_1 \oplus \mathcal{P}_2| = 1$.

Proof. We present a parameterized polynomial-time reduction from the $W[1]$ -hard EDGE-DECREMENTAL CLIQUE WITH PENDANT EDGES (see Lemma 7). Let $(G = (V, E), e^*, S^*, h)$ be an instance of EDGE-DECREMENTAL CLIQUE WITH PENDANT EDGES, where $S^* \subseteq V$ is a clique with pendant edges of size h in G . We assume without loss of generality that $|E| > \binom{h}{2} + h$.

To identify each clique vertex with its pendant edge, we introduce the following notion. Let C be a clique with pendant edges. By definition, each clique vertex from C has at least one neighbor outside C . Thus, for each clique vertex $v \in C$, let w be an arbitrary but fixed neighbor of v outside C . We use $\text{pen}_C(v)$ to denote the corresponding pendant edge $\{v, w\}$. Further, we denote by $\text{pen}(C) := \{\text{pen}(v) \mid v \in C\}$ the set of pendent edges of clique C .

In what follows, agent sets U and W will not have the same cardinality so that some agents need to remain unmatched. Note that for incomplete preferences with ties, using standard padding tricks, we can make U and W have the same cardinality and simulate an agent being unmatched by being matched with dummy agents. Since using dummy agents to “garbage collect” the unmatched agents will increase the size of symmetric difference between our initial matching and the sought matching, we need to make sure that the number of unmatched agents is upper-bounded by a function in k .

Agents, preferences, and the symmetric difference bound k . In our INCREMENTAL STABLE MARRIAGE instance, we will have one agent for each vertex from V and one agent for each edge from E . For simplicity, we use the same symbols for vertices (edges) and the corresponding agents. Further, we introduce three sets of auxiliary agents: $X := \{x_1, \dots, x_{|V|-h}\}$, $Y := \{y_1, \dots, y_{|E|-\binom{h}{2}-h-1}, y^*\}$, and $Q := \{q_1, \dots, q_t\}$, where t is an even number which is to be determined later, and two special agents y^\dagger and e^\dagger .

We partition our agents such that $U = V \cup Y \cup \{e^\dagger\} \cup \{q_1, q_3, \dots, q_{t-1}\}$ forms one part and $W = X \cup E \cup \{y^\dagger\} \cup \{q_2, q_4, \dots, q_t\}$ forms the other part. The preferences of the agents in profile \mathcal{P}_1 are set as follows.

$$\begin{aligned} \text{agent } v \in V: & \quad (\{e \in E \mid v \in e\}) \succ (X), & \text{agent } x \in X: & \quad (V) \succ q_1, \\ \text{agent } y \in Y \setminus \{y^*\}: & (E), & \text{agent } e \in E \setminus \{e^*\}: & Y \succ (\{u \in V \mid u \in e\}), \end{aligned}$$

The path agents:

$$\text{agent } q_1: \quad (X) \succ q_2, \quad \text{agent } q_t: \quad q_{t-1}$$

For each $i \in \{2, \dots, t-1\}$:

$$\text{agent } q_i: \quad q_{i-1} \succ q_{i+1},$$

Finally, we list the preferences of the four special agents associated with the extra edge $e^* = \{u^*, v^*\}$:

$$\begin{aligned} \text{agent } y^*: & \quad y^\dagger \succ E, & \text{agent } e^*: & \quad e^\dagger \succ (Y \cup \{y^*\}) \succ (\{u^*, v^*\}), \\ \text{agent } e^\dagger: & \quad y^\dagger \succ e^*, & \text{agent } y^\dagger: & \quad e^\dagger \succ y^*. \end{aligned}$$

The only agent changing its mind in profile \mathcal{P}_2 , compared to \mathcal{P}_1 , is the agent y^\dagger who switches the relative order of the (only) two potential partners in its preference list. The preference list of agent y^\dagger in \mathcal{P}_2 becomes $y^* \succ e^\dagger$. We set the difference k between the two matchings M_1 and M_2 to $k := h^2 + 5h + 4$ and set $t := 2 \cdot \lceil k/2 \rceil + 2$, which is strictly larger than k . It remains to define the matching M_1 that is stable for profile \mathcal{P}_1 to complete the construction. Before we do this, we sketch the idea of the construction.

Construction idea and the initial matching M_1 . Observe that $|W| - |U| = \binom{h}{2}$ agents will remain unmatched. As already discussed at the beginning of the proof, we can introduce $\binom{h}{2}$ dummy agents to “garbage collect” these unmatched agents. However, this will increase our parameter k only by $2 \cdot \binom{h}{2}$. To convey the actual idea of the reduction, however, we omit introducing these dummy agents.

Now, the idea of our construction is that the set $\perp(M_1)$ of agents that are not matched by our initial matching M_1 corresponds to the set of edges of the clique S^* for G . Analogously, for every stable matching M_2 for \mathcal{P}_2 the set $\perp(M_2)$ of agents that are not matched by M_2 will have to correspond to a set of edges of some clique S for $G - e^*$.

This can be formally captured as follows.

Definition 7. We define the initial matching M_1 as follows.

- (i) Match agent e^\dagger with agent y^\dagger .
- (ii) Match every clique vertex agent $v \in S^*$ with its pendant edge agent $\text{pen}_{S^*}(v)$.
- (iii) Match every non-clique vertex agent $v \in V \setminus S^*$ to an arbitrary but fixed agent from X .
- (iv) Match every agent from Y with an arbitrary but fixed non-pendant and “non-clique” edge agent $e \in E \setminus ((S^*) \cup \text{pen}(S^*))$.




Figure 3: Illustration of the stable matching M_1 for \mathcal{P}_1 and a desired stable matching M_2 for \mathcal{P}_2 (see Definition 7 and Definition 8). Observe that each agent from U is matched in both M_1 and M_2 . The agents from S and from $S^* \setminus S$ changed their partners. The agents from $\text{pen}(S)$ and from $E \setminus (\text{pen}(S) \cup \binom{S}{2}) \setminus (E \setminus (\text{pen}(S^*) \cup \binom{S^*}{2}))$ changed their partners.

(v) For every odd number $i \in \{1, 3, \dots, t-1\}$, match agent q_i with agent q_{i+1} .

This completes the construction which can be computed in polynomial time. The following claim implies that M_1 is stable for \mathcal{P}_1 . It will be used later in the correctness proof.

Claim 4. The set of unmatched agents in M_1 is $\binom{S^*}{2}$. Moreover, M_1 is stable for \mathcal{P}_1 .

Proof. The first statement follows from the fact that by Definition 7, every agent from U is matched and by Definition 7(iv), exactly those edge agents which correspond to the edges from $\binom{S^*}{2}$ are unmatched under M_1 .

As for the second statement, suppose for the sake of contradiction, that $\{u, w\}$ is a blocking pair of M_1 with $u \in U$ and $w \in W$. Now, observe that each agent from $S^* \cup (Y \setminus \{y^*\}) \cup \{e^\dagger\} \subseteq U$ already obtains its most preferred agent as a partner. Moreover, for each $i \in \{1, \dots, t/2\}$, the path agent q_{2i-1} only prefers to an agent z to its assigned partner with $z \in X \cup \{q_{2i'} \mid 1 \leq i' \leq t/2\}$, but agent z already obtains its most preferred agent as a partner (either some vertex agent or some path agent). It follows that $u \in V \setminus S^*$ or $u = y^*$. Since y^* only prefers y^\dagger to its partner but y^\dagger already obtains its most preferred agent as a partner, it follows that $u \in V \setminus S^*$, i.e., u is some non-clique vertex agent. This implies that $M_1(v) \in X$. By the preferences of the agents from V and since $M_1(u) \in X$, it follows that w is some edge agent such that $u \in w$. Since u is not a clique vertex from S^* , it follows that $w \not\subseteq \binom{S^*}{2}$, implying that $w \in E \setminus \binom{S^*}{2}$. If $w \in E \setminus ((\binom{S^*}{2} \cup \text{pen}(S^*)))$, i.e., it is also not a pendant edge of some clique vertex from S^* , then it obtains a partner from Y and will not form with u a blocking pair because u is a vertex agent. Thus, w must be a pendant edge agent from $\text{pen}(S^*)$. In this case, w is matched to another incident vertex agent which is different from u . However, since w considers both its incident vertex agents as tied, it cannot form with u a blocking pair, a contradiction. (of Claim 4) \diamond

Correctness of the Construction. We show that there is a stable matching M_2 for profile \mathcal{P}_2 with $|M_1 \Delta M_2| \leq k$ if and only if there is a clique C' with pendant edges of size h for $G - e^*$.

For the “only if” direction, let S be a clique with pendant edges pen_S of size h for $G - e^*$. For ease of notation let $E(S) = \binom{S}{2} \cup \text{pen}(S)$ and $E(S^*) = \binom{S^*}{2} \cup \text{pen}(S^*)$.

Definition 8. We construct the target matching M_2 as follows.

- (i) Match e^\dagger with e^* and y^* with y^\dagger .
- (ii) Match each clique vertex $v \in S$ with its pendant edge agent $\text{pen}_S(v)$.
- (iii) Match each non-clique vertex agent $v \in V \setminus (S \cup S^*)$ with the partner $M_1(v)$ (note that this partner is from X).
- (iv) Match each remaining non-clique vertex agent $v \in S^* \setminus S$ with an arbitrary but fixed not-yet-matched agent from X .
- (v) For each $y \in Y \setminus \{y^*\}$ with $M_1(y) \in E \setminus E(S)$, match y with $M_1(y)$; recall that $M_1(y) \in E \setminus E(S^*)$.
- (vi) Match each remaining not-yet-matched agent $y \in Y \setminus \{y^*\}$ with an arbitrary but fixed not-yet-matched agent from $E \setminus E(S)$.
- (vii) For every odd number $i \in \{1, 3, \dots, t-1\}$, match agent q_i with agent q_{i+1} .

We claim the following for matching M_2 .

Claim 5. The set of unmatched agents in M_2 is $\binom{S}{2}$. Moreover, M_2 is stable for \mathcal{P}_2 .

Proof. Since S is a clique with pendant edges for $G - e^*$, we can infer that $e^* \notin \binom{S}{2} \cup \text{pen}(S)$. Thus, M_2 is indeed a matching. Moreover, by the definition of M_2 (see Definition 8(ii), (v), (vi)) and similarly to the proof for Claim 4, we can conclude that exactly those edge agents which correspond to some edge from $\binom{S}{2}$ are unmatched under M_2 .

As for the stability proof, observe that y^\dagger 's most preferred agent in \mathcal{P}_2 is agent y^* . Using a proof similar to the one for Claim 4, we can conclude that M_2 is stable for \mathcal{P}_2 . (of Claim 5) \diamond

Now, we upper-bound the size $|M_1 \Delta M_2|$.

Claim 6. It holds that $|M_1 \Delta M_2| \leq k$.

Proof. To prove this, note that we only need to count the number of agents from U who change their partners because both M_1 and M_2 match all agents from U and unmatched exactly $\binom{h}{2}$ agents from E (see Claim 4 and Claim 5). Thus, $|M_1 \Delta M_2| \leq 2 \cdot |\{u \in U \mid M_1(u) \neq M_2(u)\}|$. Between M_1 and M_2 , the following agents from U may have changed their partners.

- (1) The clique vertex agents from S ,
- (2) the non-clique vertex agents from $S^* \setminus S$,
- (3) the agents y from $Y \setminus \{y^*\}$ with $M_2(y) \in (E \setminus E(S)) \setminus (E \setminus E(S^*))$,
- (4) the agents y^* , and e^\dagger .

Before we continue with the bound, let us recall a property in set theory.

$$\text{For each three sets } A, B, C \text{ it holds that } A \setminus (B \setminus C) = (A \cap C) \cup (A \setminus B). \quad (10)$$

Substituting $A = E \setminus E(S)$, $B = E$, and $C = E(S^*)$, we conclude the following for the set of agents described in (3).

$$(E \setminus E(S)) \setminus (E \setminus E(S^*)) = ((E \setminus E(S)) \cap E) \cup ((E \setminus E(S)) \setminus E) = (E \setminus E(S)) \cap E(S^*) \subseteq E(S^*).$$

Summarizing, by the definition of $E(S^*)$, the number of agents from U that changed their partners is at most $|S| + |S^*| + |\binom{S^*}{2}| + |\text{pen}(S^*)| + 2 = 3h + \binom{h}{2} + 2$. Thus, $|M_1 \Delta M_2| \leq 2 \cdot (3h + \binom{h}{2} + 2) = h^2 + 5h + 4$. (of Claim 6) \diamond

For the “if” direction, let M_2 be an arbitrary stable matching for \mathcal{P}_2 with $|M_1 \Delta M_2| \leq k$. Our correctness proof mainly relies on the following claim.

Claim 7. *Let $Z \subseteq V$ be the set of vertex agents that are not matched to X in M_2 and let $F = E \cap \perp(M_2)$ be the set of unmatched edge agents. The following holds.*

- (i) $e^* \notin F$.
- (ii) $|Z| = h$.
- (iii) $F = \binom{Z}{2}$.
- (iv) Every agent from Z is matched to an edge agent from $E \setminus \binom{Z}{2}$.

Proof. For the first statement, observe that by the preferences of the agents from $\{y^*, e^\dagger, y^\dagger, e^*\}$ in \mathcal{P}_2 and by the stability of M_2 for \mathcal{P}_2 it must hold that $M_2(e^\dagger) = e^*$ and $M_2(y^\dagger) = y^*$. Thus, e^* is matched, implying that $e^* \notin F$.

For the second statement, note that M_2 must match every agent from X to some agent from V as otherwise there would be an agent $x \in X$ with $M_2(x) = q_1$. Every stable matching, however, that contains $\{x, q_1\}$ must also match q_{2i} with q_{2i+1} for every number $i \in \{1, 2, \dots, t/2 - 1\}$. The symmetric difference between M_1 and M_2 is at least $2 \cdot (t/2 - 1 + 1) = t > k$ —a contradiction. Since $|X| = |V| - h$ it follows that exactly h agents from V remain that are *not* matched to X . By the definition of Z , we immediately have $|Z| = h$.

To show the third statement, first, we show that

$$F \subseteq \binom{Z}{2}. \quad (11)$$

Towards a contradiction, suppose that there is an edge $e \in F \setminus \binom{Z}{2}$, i.e., there exists an unmatched edge agent $e \in F$ for which at least one of its endpoint is matched to X . Let $v \in e$ be such an incident vertex of e with $M_2(v) \in X$. Note that by the first statement, we have that $e \neq e^*$ since e is unmatched under M_2 . Now, we claim that M_2 is not stable since $\{v, e\}$ is a blocking pair. Indeed, edge agent e is not matched and prefers to be matched with v and v finds e better than its current partner (from X). Thus, M_2 is not stable—a contradiction.

Second, we show that $|F| \geq \binom{h}{2}$. To see this, observe that by the preferences of the agents from $E \setminus \{e^*\}$, each edge agent from $E \setminus \{e^*\}$ is either unmatched or matched to a vertex agent from V or matched to an agent from $Y \setminus \{y^*\}$. Now, recall by the definition of Z that each agent from $V \setminus Z$ is matched to some agent from X . Thus, each edge agent from $E \setminus \{e^*\}$ is either unmatched or matched to an agent from $Z \cup Y \cup \{y^*\}$. By the definition of F and since $M_2(e^*) = e^\dagger$, it follows that

$$|F| \geq |E \setminus \{e^*\}| - |Z \cup (Y \setminus \{y^*\})| = |E| - 1 - (h + |E| - \binom{h}{2} - h - 1) = \binom{h}{2}.$$

The second to last equation holds because $|Z| = h$ (see the second statement). Together with (11), we infer that $|F| = \binom{h}{2}$ and thus $F = \binom{Z}{2}$.

Finally, to show the last statement, consider an arbitrary agent $z \in Z$. By the definition of Z , it follows that z is either unmatched or matched to some agent from $E \setminus F$. By the definition of F and by the third statement, there exists an unmatched edge agent $e \in F$ which is incident to z . By the stability of M_2 it must hold that z is matched to some edge agent from $E \setminus F$, i.e., from $E \setminus \binom{Z}{2}$. (of Claim 7) \diamond

We construct our clique S with pendant edges pen_S of size h for $G - e^*$ by applying Claim 7 as follows. The clique is set to $S := Z$ and for each vertex $v \in S$ we set its pendant edge $\text{pen}_S(v)$ to $M_2(v)$. Note that by Claim 7(i) it holds that $e^* \notin F$. Together with and Claim 7(iii) we infer that $|\{u^*, v^*\} \cap Z| \leq 1$ and, hence, Z is indeed a clique in $G - e^*$. By Claim 7(iv), the defined pendant edges are all from $G - e^*$. \square

In the following, we show that with respect to the number of common pairs between the target stable matching and the initial stable matching the problem is parameterized intractable, even if the two input profiles differ by only two swaps. The corresponding parameterized reduction is from INDEPENDENT SET.

Theorem 3. INCREMENTAL STABLE MARRIAGE *with ties* is $W[1]$ -hard with respect to $k' = |M_1 \cap M_2|$ of common pairs, even if $|\mathcal{P}_1 \oplus \mathcal{P}_2| = 2$.

Proof. We show this by reducing from the $W[1]$ -hard INDEPENDENT SET problem, parameterized by the solution size. The parameterized INDEPENDENT SET problem has, as input, an undirected graph G and an integer $h \in \mathbb{N}$ —the parameter—and asks whether G admits an *independent set* of size h , i.e. an h -vertex subset of $V' \subseteq V(G)$ of pairwise nonadjacent vertices. Let $I = (G, h)$ be an instance of INDEPENDENT SET. Further, let $V(G) = \{v_1, \dots, v_n\}$ and $E(G) = \{e_0, \dots, e_{m-1}\}$ denote the set of vertices and the set of edges in G respectively; note that we start the index of the edges with zero to simplify our reasoning later. We construct an instance (P_1, P_2, M_1) of INCREMENTAL STABLE MARRIAGE *with ties*, where the two profiles P_1 and P_2 are preference profiles for two disjoint sets of agents, $U \cup W \cup E \cup F \cup A \cup B$ and $X \cup Y \cup H \cup C \cup D$, and the matching M_1 is stable for P_1 such that P_1 and P_2 will differ from each other by only two swaps. We will be searching for stable matching M_2 for P_2 with $|M_1 \cap M_2| \geq 2h$.

The agent sets. For each vertex $v_i \in V(G)$, we introduce four agents u_i, w_i, x_i, y_i , and add them to U, W, X, Y , respectively.

For each edge $e_\ell \in E(G)$ with two endpoints v_i and v_j , do the following. Introduce four agents $e_\ell, f_\ell, h_\ell^{u_i}$, and $h_\ell^{u_j}$. Add e_ℓ to E , f_ℓ to F , and $h_\ell^{u_i}$ as well as $h_\ell^{u_j}$ to H . Additionally, introduce the auxiliary agents a_ℓ, b_ℓ, c_ℓ , and d_ℓ , and add them to A, B, C , and D , respectively.

The preference lists of the agents from U, W, X , and Y . The preference lists of these agents are the same in both P_1 and P_2 , and are constructed in such a way that an arbitrary stable matching must match these agents among themselves. Here, $[\star]$ means that the elements in \star are ranked in an arbitrary but fixed order, while (\star) means that the elements in \star are tied.

$$\begin{aligned} \forall i \in [n], \quad & u_i: x_i \succ [\{h_\ell^{u_i} \mid e_\ell \in E(G) \text{ with } v_i \in e_\ell\}] \succ y_i, & x_i: w_i \succ u_i, \\ & w_i: y_i \succ x_i & y_i: u_i \succ w_i. \end{aligned}$$

The preference lists of the agents from E, F , and H . The preference lists of these agents are the same in both P_1 and P_2 .

$$\begin{aligned} \forall e_\ell \in E[G] \text{ with } e_\ell = \{v_i, v_j\} \text{ and } i < j, \quad & e_\ell: (h_\ell^{u_i}, h_\ell^{u_j}) \succ c_\ell, & h_\ell^{u_i}: (e_\ell, u_i, a_\ell) \succ f_\ell, \\ & f_\ell: (h_\ell^{u_i}, h_\ell^{u_j}) \succ d_\ell, & h_\ell^{u_j}: (e_\ell, u_j, b_\ell) \succ f_\ell. \end{aligned}$$

The preference lists of the auxiliary agents from A, B, C , and D . Except for the agents c_0 and d_0 , the preference lists of these agents are the same in both P_1 and P_2 . We first describe the preference lists of all the auxiliary agents in P_1 , and then only describe the preference lists of c_0 and d_0 in P_2 as the other will remain the same. The operations “ $\ell + 1$ ” and “ $\ell - 1$ ” are taken modulo m .

$$\begin{aligned} \forall e_\ell \in E[G] \text{ with } e_\ell = \{v_i, v_j\} \text{ and } i < j, \quad & a_\ell: c_{\ell+1} \succ h_\ell^{u_i}, & c_\ell: (a_{\ell-1}, e_\ell), \\ & b_\ell: d_{\ell+1} \succ h_\ell^{u_j}, & d_\ell: (b_{\ell-1}, f_\ell). \end{aligned}$$

In P_2 , the preference lists of agents c_0 and d_0 will be changed so that the acceptable agents are not tied any more:


$$c_0: a_{m-1} \succ e_0, \text{ and } d_0: b_{m-1} \succ f_0.$$

It is straight-forward to verify that P_1 and P_2 differ by only two swaps.

The initial stable matching M_1 of P_1 . For each vertex $v_i \in V(G)$, let $M_1(u_i) = y_i$, $M_1(w_i) = x_i$. For each edge $e_\ell \in E(G)$ with $e_\ell = \{v_i, v_j\}$ and $i < j$, let $M_1(e_\ell) = c_\ell$, $M_1(f_\ell) = d_\ell$, $M_1(a_\ell) = h_\ell^{u_i}$, and $M_1(b_\ell) = h_\ell^{u_j}$. One can verify that this matching is indeed a stable matching of P_1 .

As already mentioned, we set the minimum number of common pairs between M_1 and the target matching M_2 (which shall be stable in P_2) to be $2h$. This completes the construction.

The following figure depicts a portion of acceptability graph of profile P_1 , where the labels on the edges denote the ranks of the agents. The edges marked with gray colors are part of matching M_1 .



Before we show that this is parameterized reduction, we observe the following.

Claim 8. *Every stable matching M of P_2 satisfies the following.*

1. For each vertex $v_i \in V(G)$, we have that $\{M(u_i), M(w_i)\} = \{x_i, y_i\}$.
2. For each edge $e_\ell \in E(G)$, agents c_ℓ and d_ℓ must be assigned some partners under M .
3. For each edge $e_\ell \in E(G)$ with $e_\ell = \{v_i, v_j\}$, we have that $M(c_\ell) = a_{\ell-1}$, $M(d_\ell) = b_{\ell-1}$, and $\{M(h_{\ell-1}^{u_i}), M(h_{\ell-1}^{u_j})\} = \{e_{\ell-1}, f_{\ell-1}\}$.
4. For each vertex $v_i \in V(G)$ and each edge $e_\ell \in E(G)$ with $v_i \in e_\ell$, if $M(u_i) = y_i$, then $M(h_\ell^{u_i}) = e_\ell$.

Proof. For the first statement, suppose, towards a contradiction, that there exists some vertex v_i such that $\{M(u_i), M(w_i)\} \neq \{x_i, y_i\}$. By the preference lists of x_i and y_i , it follows that x_i or y_i is not assigned a partner by M . If agent x_i was not assigned a partner, then it will form with w_i a blocking pair of M as x_i is the only agent which w_i prefers most. Analogously, if agent y_i was not assigned a partner, then it will form with u_i a blocking pair of M .

For the second statement, observe that $a_{\ell-1}$ ranks c_ℓ at the first place. If agent $c_{\ell-1}$ would have been unmatched under M , then it would form with $a_{\ell-1}$ a blocking pair. Analogously, we can obtain that $d_{\ell-1}$ must also be assigned a partner.

For the third statement, observe that in P_2 , agents c_0 and a_{m-1} form a fixed pair, meaning that they prefer each other more than any other agents. Thus, agents c_0 and a_{m-1} must be matched with each other in any stable matching of P_2 . Analogously, agents d_0 and b_{m-1} must be matched with each other in any stable matching of P_2 .

Now, let $v_{i_{m-1}}$ and $v_{j_{m-1}}$ be the two endpoints of edge e_{m-1} . By the preference lists of e_{m-1} and f_{m-1} , we know that $h_{m-1}^{u_{i_{m-1}}}$ and $h_{m-1}^{u_{j_{m-1}}}$ must be assigned some partners as otherwise the one that is unmatched (under M) will form with e_{m-1} and f_{m-1} two blocking pairs of M . By the first statement we have that $M(u_{i_{m-1}}) \in \{x_{i_{m-1}}, y_{i_{m-1}}\}$ and since a_{m-1} is already matched to c_0 , agent $h_{m-1}^{u_{i_{m-1}}}$ can only be matched to an agent that is either e_{m-1} or f_{m-1} . Analogously, agent $h_{m-1}^{u_{j_{m-1}}}$ can only be matched to an agent that is either e_{m-1} or f_{m-1} .

We have just reasoned that the partners of e_{m-1} and f_{m-1} under M are from $\{h_{m-1}^{u_{i_{m-1}}}, h_{m-1}^{u_{j_{m-1}}}\}$. By the preference lists of c_{m-1} and d_{m-1} , using the second statement, we conclude that $M(c_{m-1}) = a_{m-2}$ and $M(d_{m-1}) = b_{m-2}$. By a similar reasoning as we did for a_{m-1} and b_{m-1} , we can achieve our desired third statement.

For the fourth statement, assume that $M(u_i) = y_i$. Since u_i prefers $h_\ell^{u_i}$ to its partner y_i , by the stability of M , it follows that $h_\ell^{u_i}$ must find its partner $M(h_\ell^{u_i})$ at least as good as u_i . This means that $M(h_\ell^{u_i}) \neq f_\ell$. By the third statement, it follows that $M(h_\ell^{u_i}) = e_\ell$. (of Claim 8) \diamond

Now, we are ready to show the correctness of the construction, i.e. I admits an independent set of size h if and only if P_2 admits a stable matching M_2 with $|M_1 \cap M_2| \geq 2h$. For the “only if” direction, assume that V' is an independent set of G with h vertices. We construct matching M_2 as follows.

1. For each vertex $v_i \in V$, if $v_i \notin V'$ is *not* from the independent set, then let $M_2(u_i) = x_i$ and $M_2(w_i) = y_i$; otherwise let $M_2(u_i) = M_1(u_i) = y_i$ and $M_2(w_i) = M_1(w_i) = x_i$.
2. For each edge $e_\ell \in E(G)$, let v_i and v_j denote the endpoints of e_ℓ with $i < j$. If $v_j \in V'$ is from the independent set, implying that $v_i \notin V'$, then let $M_2(e_\ell) = h_\ell^{u_j}$ and $M_2(f_\ell) = h_\ell^{u_i}$. Otherwise, let $M_2(e_\ell) = h_\ell^{u_i}$ and $M_2(f_\ell) = h_\ell^{u_j}$. Let $M_2(a_\ell) = c_{\ell+1}$ and $M_2(b_\ell) = d_{\ell+1}$.

It is straight-forward to verify that $|M_1 \cap M_2| = 2h$ as they share at the pairs that correspond to the vertices in the independent set. Now, we focus on the stability of M_2 . Towards a contradiction, suppose that M_2 is not stable in P_2 , and let p be a blocking pair of M_2 . This pair p must involve some agent from $U \cup W \cup E \cup F \cup A \cup B$.

First of all, one can verify that no agent w_i from W would be involved in a blocking pair because x_i is the only agent with which w_i could form a blocking pair, but x_i will already obtain its most preferred agent y_i . Further, no agent from $E \cup F \cup A \cup B$ would be involved in a blocking pair as they already received one their most preferred agents. Thus, we obtain that p involves some agent from U , say u_z . We know that $M_2(u_z) = y_z$ as otherwise we have that $M_2(u_z) = x_z$ which is u_z 's most preferred agent—a contradiction to u_z being in blocking pair p . By our definition of M_2 , it follows that $v_z \in V'$ is from the independent set. Hence, the other agent in blocking pair p must be some agent $h_\ell^{u_z}$ with $v_z \in e_\ell$ such that $h_\ell^{u_z}$ prefers u_z to $M_2(h_\ell^{u_z})$. By the preference list of $h_\ell^{u_z}$ it follows that $M_2(h_\ell^{u_z}) = f_\ell$ as otherwise $h_\ell^{u_z}$ will not form with u_z a blocking pair. Let $v_{z'}$ be the other endpoint of e_ℓ , implying that $v_{z'} \notin V'$. If $z < z'$, then by our definition of M_2 , it follows that $M_2(h_\ell^{u_z}) = e_\ell$ —a contradiction. If $z' < z$, then by our definition of M_2 , it also follows that $M_2(h_\ell^{u_z}) = e_\ell$ —a contradiction. This shows that M_2 is indeed a stable matching of P_2 which shares with M_1 by $2h$ pairs.

For the other direction, assume that M_2 is a stable matching of P_2 , sharing with M_1 by at least $2h$ pairs. By the third statement in Claim 8, it follows that M_2 can only share with M_1 by the pairs that involve some agents from $U \cup W \cup X \cup Y$. Now we construct a vertex subset according to these pairs. Let $V' = \{v_i \in V \mid M_2(u_i) = y_i\}$. Clearly, using the first statement in Claim 8, we have that $|V'| \geq h$ as $|M_1 \cap M_2| \geq 2h$. Now, to show that V' is indeed an independent set, suppose, towards a contradiction, that V' has two adjacent vertices, denoted as v_i and v_j . Let e_ℓ be the incident edge of v_i and v_j . By the fourth statement of Claim 8 we have that $M_2(h_\ell^{u_i}) = e_\ell = M_2(h_\ell^{u_j})$ —a contradiction to M_2 being a matching. \square

Finally, we show that even a single swap in the preference list of one single agent makes the INCREMENTAL STABLE ROOMMATES problem $W[1]$ -hard with respect to k when ties are allowed. To show this result, we give a reduction from EDGE-INCREMENTAL INDEPENDENT SET. The construction idea is inspired by a reduction from VERTEX COVER to STABLE ROOMMATES with structured preferences [5], which, however, is relying on incomplete preferences and not showing parameterized intractability.

Theorem 4. INCREMENTAL STABLE ROOMMATES *with ties and complete preferences* is $W[1]$ -hard with respect to $k = |M_1 \Delta M_2|$, even if $|\mathcal{P}_1 \oplus \mathcal{P}_2| = 1$. It remains NP-hard even if $|\mathcal{P}_1 \oplus \mathcal{P}_2| = 1$, and the sought stable matching M_2 only needs to satisfy that $|M_1 \cap M_2| \geq 0$.




Figure 4: An illustration of the hardness reduction for Theorem 4.

Proof. Let $(G = (V, E), e^* \in E, h, S^* \subseteq V)$ be an EDGE-INCREMENTAL INDEPENDENT SET instance with $r := |V|$ vertices. We assume w.l.o.g. that $h < r$, $S^* = \{v_{r-h+1}, \dots, v_r\}$. Moreover, we assume that $e^* \subseteq S^*$ (as otherwise the input instance is a trivial yes-instance) with $e^* = \{v_{r-1}, v_r\}$. We construct an INCREMENTAL STABLE ROOMMATES instance $(\mathcal{P}_1, \mathcal{P}_2, M_1)$ with agent set U and $|\mathcal{P}_1 \oplus \mathcal{P}_2| = 1$. We will show that G has an independent of size at least h if and only if \mathcal{P}_2 admits a stable matching M_2 with $|M_1 \oplus M_2| \leq 4h$.

Before we describe the construction, we prove the following claim which is heavily used in our preference profile construction to force two agents to be matched together.

Claim 9. *Let \mathcal{P} be a profile for an agent set U , and let x, y , and z be three distinct agents with the following preference lists,*

$$\begin{aligned} \text{agent } x: & V \succ y \succ z \succ \dots, \\ \text{agent } y: & z \succ x \succ \dots, \\ \text{agent } z: & x \succ y \succ \dots, \end{aligned}$$

where $V \subseteq U \setminus \{x, y, z\}$ is a non-empty subset of agents, The symbol “...” at the preference list of each agent a denotes an arbitrary but fixed order of all remaining agents, other than a and not explicitly stated before “...”: Then, every stable matching M for \mathcal{P} must fulfill that (i) $M(x) \in V$ and (ii) $\{y, z\} \in M$.

Proof. Assume towards a contradiction to (i) that \mathcal{P} admits a stable matching M with $M(x) \notin V$. There are three cases: (1) $M(x) = y$, implying the blocking pair $\{y, z\}$, (2) $M(x) = z$, implying the blocking pair $\{x, y\}$, and (3) $M(x) \notin \{y, z\} \cup V$, implying the blocking pair $\{x, z\}$. Thus, x must be matched with some agent from V . For (ii), statement (i) implies that neither y nor z is matched with x . Now, if $\{y, z\} \notin M$, then $\{y, z\}$ forms a blocking pair. (of Claim 9) \diamond

Main idea and the constructed agents. To explain the main idea of the reduction, we first describe the agent set U and the corresponding non-complete acceptability graph of \mathcal{P} as illustrated through an example in Figure 4. At the end of the proof, we show that we can adjust the reduction so that the acceptability graph becomes complete.

For each vertex $v_i \in V$, we introduce a *vertex agent* v_i (for the sake of simplicity, we use the same symbol for the vertex and the corresponding agent). Additionally, there is a set of *cover agents* $C := \{c_1, \dots, c_{r-h}\}$ as well as three sets of *selector agents* $X := \{x_1, x_2, \dots, x_h\}$, $Y := \{y_1, y_2, \dots, y_h\}$, and $Z := \{z_1, z_2, \dots, z_h\}$. The agent set U is defined as $V \cup C \cup X \cup Y \cup Z$. For the acceptability graph, we have that every vertex agent v_i accepts every cover agent from C , every selector agent from X , and every vertex agent v_j that corresponds to a neighbor of v_i in the input graph G . For each $i \in \{1, 2, \dots, h\}$, the collector agents x_i, y_i , and z_i pairwise accept each other.

Using [Claim 9](#) we construct the preference profile \mathcal{P}_2 of the agents such that in every stable matching only the cover agents from C and the selector agents from X can be matched to the vertex agents and the vertex agents matched to the selector agents from X correspond to an independent set (of size $|X| = h$). These properties are given by the subsequent [Claim 11](#).

Agent preferences in \mathcal{P}_1 and \mathcal{P}_2 . First, we describe the preferences profiles \mathcal{P}_2 that realize the idea and the acceptability graph as described above.

$$\begin{aligned}
\mathcal{P}_2: \text{agent } v_i: [C] \succ [N(v_i)] \succ x_1 \succ x_2 \succ \dots \succ x_h \succ \dots, & \quad \forall 1 \leq i \leq r-1, \\
\text{agent } v_r: [C] \succ [N(v_r) \setminus \{v_{r-1}\}] \succ v_{r-1} \succ x_1 \succ x_2 \succ \dots \succ x_h \succ \dots, & \\
\text{agent } c_i: v_1 \sim v_2 \sim \dots \sim v_r \succ \dots, & \quad \forall 1 \leq i \leq r-h, \\
\text{agent } x_i: v_1 \sim v_2 \sim \dots \sim v_r \succ y_i \succ z_i \succ \dots, & \quad \forall 1 \leq i \leq h, \\
\text{agent } y_i: z_i \succ x_i \succ \dots, & \quad \forall 1 \leq i \leq h, \\
\text{agent } z_i: x_i \succ y_i \succ \dots, & \quad \forall 1 \leq i \leq h.
\end{aligned}$$

With a single swap in the preference list of agent v_r , we obtain the profile \mathcal{P}_1 :

$$\begin{aligned}
\mathcal{P}_1: \text{agent } v_i: [C] \succ [N(v_i)] \succ x_1 \succ x_2 \succ \dots \succ x_h \succ \dots, & \quad \forall 1 \leq i \leq r-1, \\
\text{agent } v_r: [C] \succ [N(v_r) \setminus \{v_{r-1}\}] \succ x_1 \succ v_{r-1} \succ x_2 \succ \dots \succ x_h \succ \dots, & \\
\text{agent } c_i: v_1 \sim v_2 \sim \dots \sim v_r \succ \dots, & \quad \forall 1 \leq i \leq r-h, \\
\text{agent } x_i: v_1 \sim v_2 \sim \dots \sim v_r \succ y_i \succ z_i \succ \dots, & \quad \forall 1 \leq i \leq h, \\
\text{agent } y_i: z_i \succ x_i \succ \dots, & \quad \forall 1 \leq i \leq h, \\
\text{agent } z_i: x_i \succ y_i \succ \dots, & \quad \forall 1 \leq i \leq h.
\end{aligned}$$

Herein, for each agent subset D , we denote by $[D]$ the lexicographic order of the agents by their names (resp. by their indices), called the *canonical order*. The symbol “...” at the preference list of each agent a denotes an arbitrary but fixed order of all remaining agents, other than a and not explicitly stated before “...”:

Stable matching M_1 for \mathcal{P}_1 . The initial matching M_1 is defined as follows:

- for each $i \in \{1, 2, \dots, h\}$ set $M(v_{r-i+1}) := x_i$;
- for each $i \in \{1, 2, \dots, r-h\}$ set $M(v_i) = c_i$;
- for each $i \in \{1, 2, \dots, h\}$ set $M(y_i) = z_i$.

This completes the construction and can clearly be performed in polynomial time.

Correctness of the construction. First of all, we claim that in the constructed INCREMENTAL STABLE ROOMMATES instance M_1 is indeed a stable matching for \mathcal{P}_1 .

Claim 10. *The matching M_1 is a stable matching for the profile \mathcal{P}_1 .*

Proof of Claim 10. First, observe that, by definition of M_1 , no agent from C or X can be involved in a blocking pair, since they all are matched to one of their (tied) favorite partners. Second, since the agents from X are not available also the agents from Y and from Z cannot be involved in blocking pairs. Finally, a blocking pair that consist of two agents from V (the last possibility for a blocking pair to emerge) would mean that we have two agents v_{r-i+1} and v_{r-j+1} ($1 \leq i \neq j \leq h$) that are both matched to some agent from X . Furthermore, in order to be a blocking pair both agents must be adjacent in G because only agents corresponding to neighbors in G are preferred to agents from X . Since the vertices v_{r-h+1}, \dots, v_r are an independent set in

$G - e^*$ for $e^* = \{v_{r-1}, v_r\}$, only $\{v_{r-1}, v_r\}$ could be a blocking pair. However, $\{v_{r-1}, v_r\}$ is also not blocking M_1 because (due to the swap of x_1 and v_{r-1} in the preferences of v_r between \mathcal{P}_2 and \mathcal{P}_1) agent v_r prefers its partner $M_1(v_r) = x_1$ over v_{r-1} . (of Claim 10) \diamond

Second, we claim the following:

Claim 11. *Every stable matching M'_2 for \mathcal{P}_2 satisfies the following two properties:*

1. *every vertex agent v_i is matched to either a cover agent from C or a selector agent from X , that is, $M(v_i) \in C \cup X$ for every $v \in V$, and*
2. *no two vertex agents that are both matched to the selector agents are adjacent in G .*

Proof of Claim 11. Let M'_2 be a stable matching for \mathcal{P} . For the first statement, observe Claim 9 immediately implies that for every selector agent $x_i \in X$, it holds that $M(x_i) \in V$. Thus, there are exactly $r - h$ vertex agents left that are not matched to agents from X . Suppose towards a contradiction that some cover agent c_j is not matched to any vertex agent, implying that at least one vertex agent v_i is left with $M(v_i) \notin X \cup C$. This, however, implies that $\{c_j, v_i\}$ is a blocking pair for M'_2 —a contradiction. For the second statement, suppose towards a contradiction that there are two vertex agents v_i, v_j with $\{M(v_i), M(v_j)\} \subseteq X$ as well as $\{v_i, v_j\} \in E$. The preference lists of v_i and v_j immediately imply that agents v_i and v_j form a blocking pair—a contradiction. (of Claim 11) \diamond

Now, we show that G has an independent set of size at least h if and only if \mathcal{P}_2 admits a stable matching M_2 with $|M_1 \oplus M_2| \leq 4h$.

The “if” part follows immediately from Claim 11 because it directly implies that the vertices corresponding to the h vertex agents matched to selector agents from X are pairwise non-adjacent in the graph G .

For the “only if” part, assume that $S \subseteq V$ be an independent set of size h for G . Let $S := \{v_{s_1}, v_{s_2}, \dots, v_{s_h}\}$ with $s_1 < s_2 < \dots < s_h$ such that the canonical order \succ implies $v_{s_1} \succ v_{s_2} \succ \dots \succ v_{s_h}$. We construct a matching M_2 (based on M_1) such that $|M_1 \oplus M_2| \leq 4h$ and such that M_2 is stable for \mathcal{P}_2 . First, we match all selector agents and the vertex agents corresponding to the independent set S :

- for each $i \in \{1, 2, \dots, h\}$ set $M_2(y_i) := z_i (= M_1(y_i))$;
- for each $i \in \{1, 2, \dots, h\}$ set $M_2(v_{s_i}) := x_i$.

So far, at most $2h$ agents have different partners in M_1 and M_2 . At least $r - 2h$ vertices are neither in the independent set S for G nor in the independent set $S^* = \{v_{r-h+1}, \dots, v_r\}$ for $G - e^*$. We keep the partners for the corresponding vertex agents unchanged:

- for each $v \in V \setminus (S \cup S^*)$ set $M_2(v) := M_1(v)$.

Finally, some number $q \leq h$ of vertex agents, namely those corresponding to vertices from $S^* \setminus S$, are so far unmatched in M_2 . Symmetrically, also $q \leq h$ cover agents, namely those cover agents that were matched to vertex agents from $S^* \setminus S$ in M_1 , are also so far unmatched in M_2 . We pair these agents arbitrarily such that each vertex agent is matched to a cover agent to finalize the definition of M_2 so far. Doing this, at most $2q \leq 2h$ agents will get different partners in M_2 when compared with M_1 . Hence, $|M_1 \Delta M_2| \leq 4h$.

It remains to show that M_2 is stable. This can be done analogously to the proof of Claim 10. First, observe that, by the definition of M_2 , no agent from C or X can be involved in a blocking pair since they all are matched to one of their (tied) favorite partners. Second, since agents from X are not available, the agents from Y and from Z cannot be involved in blocking pairs. Finally, a blocking pair that consist of two agents from V (the last possibility for a blocking pair to emerge) would mean that we have two agents v_i and v_j that are both matched to some

agent from X . Furthermore, in order to be a blocking pair both agents must be adjacent in G because only agents corresponding to neighbors in G are preferred to agents from X . This is a contradiction to our definition of M_2 , where the h agents from X have been matched to h vertex agents corresponding to the independent set S .

Unbounded k . Note that in this proof, we never used an explicit bound on the parameter k for any proof argument to work. In particular, if there is any stable matching for \mathcal{P}_2 , then there is one with distance $|M_1 \oplus M_2| \leq 4h$ (this follows from [Claim 11](#) and from the idea given in the “only if” part). In particular, the problem remains NP-hard if we set $k = 2n$, that is, $k' = 2n - k = 0$. In other words, the problem remains hard even if the distance between the matchings M_1 and M_2 is unbounded. \square

5 Conclusion

Motivated by dynamically changing preferences and the necessity to adapt the corresponding solutions, we introduced an “incremental view” on the computation of stable matchings. We believe that there are plenty of opportunities for future research, including, for instance, to study the role of parameters measuring the number of ties—in many hardness reductions this parameter is unbounded. We also left open whether INCREMENTAL STABLE ROOMMATES without ties is fixed-parameter tractable when parameterized by the swap distance between the two input preference profiles. Naturally, there are also future directions concerning more conceptual work, e.g., also studying further stability concepts in the context of our incremental model.

References

- [1] F. N. Abu-Khzam, J. Egan, M. R. Fellows, F. A. Rosamond, and P. Shaw. On the parameterized complexity of dynamic problems. *Theoretical Computer Science*, 607:426–434, 2015. [3](#)
- [2] H. Aziz, P. Biró, S. Gaspers, R. de Haan, N. Mattei, and B. Rastegari. Stable matching with uncertain linear preferences. In *Proceedings of the 9th International Symposium on Algorithmic Game Theory*, pages 195–206, 2016. [3](#)
- [3] S. Bhattacharya, M. Hoefer, C. Huang, T. Kavitha, and L. Wagner. Maintaining near-popular matchings. In *Proceedings of the 42nd International Colloquium on Automata, Languages, and Programming*, pages 504–515, 2015. [2](#), [3](#)
- [4] H. Böckenhauer, E. Burjons, M. Raszyk, and P. Rossmanith. Reoptimization of parameterized problems. *CoRR*, abs/1809.10578, 2018. [3](#)
- [5] R. Brederbeck, J. Chen, U. P. Finnendahl, and R. Niedermeier. Stable roommate with narcissistic, single-peaked, and single-crossing preferences. In *Proceedings of the 5th International Conference on Algorithmic Decision Theory*, pages 315–330, 2017. [26](#)
- [6] M. Charikar, C. Chekuri, T. Feder, and R. Motwani. Incremental clustering and dynamic information retrieval. *SIAM Journal on Computing*, 33(6):1417–1440, 2004. [1](#), [3](#)
- [7] J. Chen, D. Hermelin, M. Sorge, and H. Yedidsion. How hard is it to satisfy (almost) all roommates? In *Proceedings of the 45th International Colloquium on Automata, Languages, and Programming*, pages 35:1–35:15, 2018. [18](#)
- [8] J. Chen, R. Niedermeier, and P. Skowron. Stable marriage with multi-modal preferences. In *Proceedings of the 19th ACM Conference on Economics and Computation*, pages 269–286, 2018. [3](#)

- [9] J. Chen, P. Skowron, and M. Sorge. Matchings under preferences: Strength of stability and trade-offs. In *Proceedings of the 20th ACM Conference on Economics and Computation*, pages 41–59, 2019. 3
- [10] Á. Cseh and D. F. Manlove. Stable marriage and roommates problems with restricted edges: Complexity and approximability. *Discrete Optimization*, 20:62–89, 2016. 3
- [11] J. Drummond and C. Boutilier. Elicitation and approximately stable matching with partial preferences. In *Proceedings of the 23rd International Joint Conference on Artificial Intelligence*, pages 97–105, 2013. 3
- [12] T. Feder. A new fixed point approach for stable networks and stable marriages. *Journal of Computer and System Sciences*, 45(2):233–284, 1992. 18, 19
- [13] B. Genc, M. Siala, B. O’Sullivan, and G. Simonin. Finding robust solutions to stable marriage. In *Proceedings of the 26th International Joint Conference on Artificial Intelligence*, pages 631–637, 2017. 3
- [14] B. Genc, M. Siala, G. Simonin, and B. O’Sullivan. On the complexity of robust stable marriage. In *Proc. COCOA-17*, pages 441–448, 2017. 3
- [15] B. Genc, M. Siala, G. Simonin, and B. O’Sullivan. Robust stable marriage. In *Proceedings of the 31st AAAI Conference on Artificial Intelligence*, pages 4925–4926, 2017. 3
- [16] B. Genc, M. Siala, G. Simonin, and B. O’Sullivan. Complexity study for the robust stable marriage problem. *Theoretical Computer Science*, 775:76–92, 2019. 3
- [17] P. Ghosal, A. Kunysz, and K. E. Paluch. The dynamics of rank-maximal and popular matchings. *CoRR*, abs/1703.10594, 2017. 2, 3
- [18] D. Gusfield. The structure of the stable roommate problem: Efficient representation and enumeration of all stable assignments. *SIAM Journal on Computing*, 17(4):742–769, 1988. 4, 6, 7, 8, 9, 10, 12, 13, 16
- [19] D. Gusfield and R. W. Irving. *The Stable Marriage Problem—Structure and Algorithms*. Foundations of Computing Series. MIT Press, 1989. 5, 6, 8, 9, 33, 34, 35
- [20] S. Hartung and R. Niedermeier. Incremental list coloring of graphs, parameterized by conservation. *Theoretical Computer Science*, 494:86–98, 2013. 3
- [21] R. W. Irving. An efficient algorithm for the stable roommates problem. *Journal of Algorithms*, 6(4):577–595, 1985. 6, 7
- [22] R. W. Irving. Stable marriage and indifference. *Discrete Applied Mathematics*, 48(3):261–272, 1994. 4, 8
- [23] V. Kanade, N. Leonardos, and F. Magniez. Stable matching with evolving preferences. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*, volume 60 of *LIPICs*, pages 36:1–36:13, 2016. ISBN 978-3-95977-018-7. 3
- [24] R. Krithika, A. Sahu, and P. Tale. Dynamic parameterized problems. *Algorithmica*, 80(9):2637–2655, 2018. 3
- [25] J. Luo, H. Molter, A. Nichterlein, and R. Niedermeier. Parameterized dynamic cluster editing. In *Proceedings of the 38th International Conference on Foundations of Software Technology and Theoretical Computer Science*, pages 46:1–46:15, 2018. 3

- [26] T. Mai and V. V. Vazirani. Finding stable matchings that are robust to errors in the input. In *Proceedings of the 26th Annual European Symposium on Algorithms*, pages 60:1–60:11, 2018. 3
- [27] T. Mai and V. V. Vazirani. A generalization of Birkhoff’s theorem for distributive lattices, with applications to robust stable matchings. Technical report, arXiv:1804.05537 [cs.DM], 2018. 3
- [28] D. F. Manlove. *Algorithmics of Matching Under Preferences*. WorldScientific, 2013. 2, 5, 9
- [29] D. Marx and I. Schlotter. Parameterized complexity and local search approaches for the stable marriage problem with ties. *Algorithmica*, 58(1):170–187, 2010. 3
- [30] D. Marx and I. Schlotter. Stable assignment with couples: Parameterized complexity and local search. *Discrete Optimization*, 8(1):25–40, 2011. 3
- [31] S. Miyazaki and K. Okamoto. Jointly stable matchings. In *Proceedings of the 28th International Symposium on Algorithms and Computation*, pages 56:1–56:12, 2017. 3
- [32] P. Nimbhorkar and V. Rameshwar. Dynamic rank-maximal and popular matchings. *Journal of Combinatorial Optimization*, 37(2):523–545, 2019. 2, 3
- [33] B. Schieber, H. Shachnai, G. Tamir, and T. Tamir. A theory and algorithms for combinatorial reoptimization. *Algorithmica*, 80(2):576–607, 2018. 3

A Incremental Stable Marriage without ties

In this section, we show that INCREMENTAL STABLE MARRIAGE without ties can be solved in polynomial time, using an idea similar to the one for finding the so-called *maximum weight stable matching*, a stable matching whose corresponding closed subset of rotations have maximum weight [19, Chapter 3.6.2]; the concept revolving around the rotations will be defined shortly. Gusfield and Irving [19] presented a network flow approach to find such a maximum-weight stable matching in $O(n^2 \cdot w)$ time, where n and w denote the number of agents and the sum of weights of rotations, respectively. We will show that for our problem the sum of the weights of rotations is bounded by n , which is the size of the intersection between our sought stable matching and the target stable matching.

In the remainder of the section, we present necessary notions and the definition of the weights of the rotations and refer to Chapter 3.6.1 by Gusfield and Irving for further details.

A.1 Preliminaries

The approach will heavily utilize the structural properties revolving around stable matchings and rotations of a STABLE MARRIAGE instance without ties.

As already observed in the literature, for each STABLE MARRIAGE instance P with two disjoint sets U and W of agents, when operating on the agent set U , the Gale-Shapely algorithm always returns a U -optimal stable matching M . The matching M is *U -optimal stable matching* if it is stable and there is no other stable matching M' such that an agent from U would prefer its partner from M to the partner from M' . Starting from the U -optimal stable matching, we can successively eliminate the so-called rotations to obtain further stable matchings.

Definition 9 (Successor agent, rotations, and rotation elimination). *Let P be a STABLE MARRIAGE instance with two disjoint sets of agents, U and W , and with (possibly) incomplete preferences. Given a stable matching M for P , for each agent $u \in U$, we define its successor $\text{succ}_M(u)$ as the first (after $M(u)$) agent w on the preference list of u such that w is matched under M and prefers u to its partner $M(w)$.*

A sequence $\rho = ((u_0, w_0), (u_1, w_1), \dots, (u_{r-1}, w_{r-1}))$ of pairs is called a rotation if there exists a stable matching M for P such that for each $i \in \{0, 1, \dots, r-1\}$ we have $(u_i, w_i) \in U \times W$, $M(u_i) = w_i$, and $\text{succ}_M(u_i) = w_{i+1}$ (index $i+1$ taken modulo r). We say that rotation ρ is exposed in M .

We use the notation M/ρ to refer to the matching resulting from M by replacing each pair $\{u_i, w_i\}$ with $\{u_i, w_{i+1}\}$. Formally,

$$M/\rho = M \setminus \{\{u_i, w_i\} \mid 0 \leq i \leq r-1\} \cup \{\{u_i, w_{i+1}\} \mid 0 \leq i \leq r-1\}.$$

The transformation of M to M/ρ is called the elimination of ρ from M .

We illustrate the concept of the successor below:

$$u: \dots M(u) \dots \text{succ}_M(u) \dots, \quad \text{succ}_M(u): \dots u \dots M(\text{succ}_M(u)) \dots$$

Eliminating a rotation from a stable matching results in another stable matching [19, Lemma 2.5.2].

Definition 10 (Predecessors of rotations, the rotation poset, and the rotation digraph). *Let π and ρ be two rotations for a STABLE MARRIAGE instance P . We say that π is a predecessor of ρ , written as $\pi \triangleright^P \rho$, if no stable matching in which ρ is exposed can be obtained from the U -optimal stable matching by a sequence of eliminations of rotations without eliminating π first. The reflexive closure of the relation \triangleright^P , denoted as \supseteq^P , defines a partial order on the set of all rotations and is called the rotation poset for P . We abbreviate the name of a subset of the poset that is closed under predecessors as a closed subset.*

An alternative representation of the rotation poset $\triangleright(P)$ is through an acyclic directed graph, called rotation digraph of P and written as $G(P)$, whose vertex set is the set of rotations of P , and there is a direct arc from rotation π to rotation ρ if and only if π precedes ρ and there is no other rotation σ such that $\pi \triangleright^P \sigma \triangleright^P \rho$.

Finally, let us describe a central result from the literature that relates rotations and stable matchings.

Proposition 7 ([19, Theorem 2.5.7, Lemma 3.3.2]). *Let R denote the set of all rotations of a preference profile P , and let $G(P)$ denote the rotation digraph of P . The following holds.*

1. *A matching M is a stable matching of P if and only if there is a closed subset of rotations $R' \subseteq R$ with respect to the precedence relation \triangleright^P such that M can be generated by taking the U -optimal stable matching and by eliminating the rotations in R' in an order consistent with \triangleright^P .*
2. *The rotation set R and the rotation digraph $G(P)$ can be computed in $O(n^2)$ time.*

A.2 From Incremental Stable Marriage to Finding Maximum-Weight Closed Subset of Rotations

Given an instance $I = (\mathcal{P}_1, \mathcal{P}_2, M_1, k)$ of INCREMENTAL STABLE MARRIAGE, let M_0 be the U -optimal stable matching of \mathcal{P}_2 , let R_2 denote the set of rotations for \mathcal{P}_2 , and let G_2 be the rotation digraph for \mathcal{P}_2 . Towards finding a stable matching which shall be as close to the input matching M_1 as possible, we assign a weight to each rotation from R_2 which shall indicate the benefit of eliminating this rotation.

For each rotation $\rho \in R_2$ with $\rho = ((x_0, y_0), \dots, (x_{r-1}, y_{r-1}))$ let

$$w(\rho) := |\{(x_i, y_{i+1}) \mid \{x_i, y_{i+1}\} \in M_1, 0 \leq i \leq r-1\}| - |\{(x_i, y_i) \mid \{x_i, y_i\} \in M_1, 0 \leq i \leq r-1\}|.$$

That is, we count the number of pairs in M_1 the elimination of ρ introduces minus the number of pairs in M_1 we loose when eliminating ρ . By the above definition, we can derive the following.

Lemma 8. *If $\rho \in R_2$ is a rotation exposed in a stable matching M for \mathcal{P}_2 , then*

$$|M_1 \cap (M/\rho)| = |M_1 \cap M| + w(\rho).$$

Proof. Let $\rho = ((x_0, y_0), \dots, (x_{r-1}, y_{r-1}))$ be a rotation exposed in the stable matching M . In the following, all subscripts $i+1$ are taken modulo r . By the definition of M/ρ we have that $M/\rho = \{\{x, y\} \in M \mid (x, y) \notin \rho\} \uplus \{\{x_i, y_{i+1}\} \mid 0 \leq i \leq r-1\}$. Thus, we prove the statement by showing the following; note that no $\{x_i, y_i\}$ belongs to M/ρ , $0 \leq i \leq r-1$.

$$\begin{aligned} |M_1 \cap (M/\rho)| &= |M_1 \cap ((M/\rho) \cap M)| + |M_1 \cap ((M/\rho) \setminus M)| \\ &= |\{\{x, y\} \in M_1 \mid \{x, y\} \in M \wedge (x, y) \notin \rho\}| + \\ &\quad |\{\{x, y\} \in M_1 \mid (x, y) = (x_i, y_{i+1}) \text{ for some } 0 \leq i \leq r-1\}| \\ &= |\{\{x, y\} \in M_1 \cap M\}| - |\{\{x, y\} \in M_1 \cap M \mid (x, y) \in \rho\}| + \\ &\quad |\{\{x, y\} \in M_1 \mid (x, y) = (x_i, y_{i+1}) \text{ for some } 0 \leq i \leq r-1\}| \\ &= |M_1 \cap M| + w(\rho). \end{aligned}$$

Note that the second to last equation holds because $(x, y) \in \rho$ implies that $\{x, y\} \in M$. \square

By applying Lemma 8 repeatedly, we obtain the following.

Corollary 1. *If $C \subseteq R_2$ is the (unique) closed subset of rotations associated with stable matching M of \mathcal{P}_2 , then $|M_1 \cap M| = |M_1 \cap M_0| + \sum_{\rho \in C} w(\rho)$.*

Since a stable matching M_2 for \mathcal{P}_2 with minimum symmetric difference $\text{dist}(M_1, M_2)$ has the maximum intersection $M_1 \cap M_2$, we obtain the following.

Lemma 9. *Let M_2 be a stable matching for \mathcal{P}_2 and let C be the associated closed subset of rotations. Then, $\text{dist}(M_1, M_2) \leq k$ if and only if $\sum_{\rho \in C} w(\rho) \geq \frac{\text{dist}(M_1, M_0) - k}{2}$.*

Proof. By the definition of symmetric difference, we derive the following.

$$\text{dist}(M_1, M_2) = |M_1| + |M_2| - 2|M_1 \cap M_2|. \quad (12)$$

Since for preferences without ties, all stable matchings match the same set of agents (Proposition 1), we have that $|M_0| = |M_2|$. Together with Corollary 1, (12) is equivalent to

$$\begin{aligned} \text{dist}(M_1, M_2) &= |M_1| + |M_0| - 2 \left(|M_1 \cap M_0| + \sum_{\rho \in C} w(\rho) \right) \\ &= \text{dist}(M_1, M_0) - 2 \sum_{\rho \in C} w(\rho). \end{aligned}$$

Our statement follows immediately. \square

By the above, our problem reduces to finding a maximum-weight closed subset of rotations. In the following, we show that the sum of the weights of the rotations is at most n and finding such a subset of rotations can thus be done efficiently.

Lemma 10. $\sum_{\rho \in R_2} w(\rho) \leq |M_1|$ and finding a closed subset of rotations with maximum weight can be done in $O(n^3)$ time.

Proof. Slightly abusing the intersection notation, for each rotation $\rho = ((x_0, y_0), \dots, (x_{r-1}, y_{r-1})) \in R_2$ we define $\rho \cap M_1 := \{(x_i, y_{i+1}) \mid \{x_i, y_{i+1}\} \in M_1\}$ ($i + 1$ is taken modulo r).

Next, observe that by one of the statement in [19, Lemma 3.2.1] implies that for each pair (x, y) there exists at most one rotation ρ such that $(x, y) \in \rho \cap M_1$. Moreover, by definition, we have that $w(\rho) \leq |\rho \cap M_1|$. Summarizing, we have that

$$\sum_{\rho \in R_2} w(\rho) \leq \sum_{\rho \in R_2} |\rho \cap M_1| = \left| \bigcup_{\rho \in R_2} (\rho \cap M_1) \right| \leq |M_1|.$$

Gusfield and Irving [19, Theorem 3.6.2] showed that finding a maximum-weight closed subset of rotations can be reduced to finding a minimum s - t cut in a specific flow network which features the precedence relation of the rotations, where the numbers of vertices and arcs are in $O(n^2)$ and the minimum cost of an s - t cut is bounded by the sum of the weights of the rotations. The latter problem can be solved in $O(|E| \cdot w)$ time (by using Ford-Fulkerson's algorithm), where $|E|$ denotes the number of arcs in the network and w is the cost of the minimum s - t cut. Since $|E| \in O(n^2)$ and since $w \leq \sum_{\rho \in R_2} w(\rho)$, we can find a maximum-weight closed subset of rotations in $O(n^2 \cdot |M_1|) = O(n^3)$ time. \square